



Ultrasound-guided bilateral greater occipital nerve block on headache seen after endovascular treatment of ruptured or unruptured intracranial aneurysms: A case report

Rüptüre olmuş ve olmamış intrakraniyal anevrizmaların endovasküler tedavisi sonrası görülen baş ağrısında ultrason eşliğinde bilateral büyük oksipital sinir bloğu: Bir olgu sunumu

▯ Rafi DOĞAN, ▯ Hüseyin Ulaş PINAR, ▯ Ömer KARACA, ▯ Fatma KARAKOÇ

Summary

Headache is a common symptom in subarachnoid hemorrhage (SAH). Often, pain control is difficult and the patient's consciousness becomes more complicated with opioids. The study aimed to reduce opioid consumption and obtain effective pain control on headache seen after endovascular treatment of intracranial aneurysms through ultrasound-guided bilateral greater occipital nerve (GON) block. Case 1 – A 59-year-old male patient with a Glasgow Coma Scale (GCS) of 13 was diagnosed with Fisher 3 grade SAH; coiling and stenting was performed for anterior communicating artery aneurysm. CSF drainage was provided with lumbar spinal catheter. Case 2 – A 55-year-old male patient with GCS of 15 who underwent coiling of the fusiform aneurysm in the left basilar artery and stenting of the stenotic region due to the basilar artery aneurysm was evaluated. Visual analog scores (VAS) of patients were 9 and 7 respectively, bilateral greater occipital nerve block with ultrasound guidance was performed. The VAS scores of both patients decreased to 3. VAS did not exceed 3 during the follow-up in intensive care unit, so there was no need for additional analgesics. In conclusion, bilateral GON block provided effective analgesia and significantly reduced additional analgesic need in both cases.

Keywords: Aneurysmal subarachnoid hemorrhage; headache; occipital nerve blockade.

Özet

SAK olgularında baş ağrısı sık görülen semptomdur. Çoğu zaman ağrı kontrolü güç olmakta ve opioid kullanılarak hastanın şuur durumu daha karmaşık hale gelmektedir. Çalışmada, ultrason eşliğinde bilateral büyük oksipital sinir bloğu ile intrakraniyal anevrizmaların endovasküler tedavisi (EVT) sonrası görülen baş ağrısının etkili kontrolünü ve opioid tüketimini azaltmayı amaçladık. Olgu1 – Fisher 3 grade SAK tanısıyla gelen, GKS: 13 olan 59 yaşında erkek hastaya, anterior kominikan arterde anevrizma nedeniyle koilizasyon ve stent uygulaması yapıldı. Lomber spinal kateterle BOS drenajı sağlandı. Olgu 2 – Baziller arter anevrizması nedeniyle, sol baziller arterde fusiform anevrizmaya koilizasyon ve stenotik bölgeye stent uygulanan GKS: 15 olan 55 yaş erkek hasta değerlendirildi. VAS skorları sırasıyla 9–7 olan hastalara, ultrason eşliğinde bilateral büyük oksipital sinir bloğu sağlandı. Her iki hastanın VAS skoru 3 ün altına düştü. Skor takip boyunca 3'ü aşmadı ve ek analjezik ihtiyacı olmadı. Sonuç olarak, iki olguda da bilateral oksipital sinir bloğu ile etkili analjezi sağlanmıştır ve ek analjezik ihtiyacı çok azalmıştır.

Anahtar sözcükler: Anevrizmal subarahnoid kanama; baş ağrısı; oksipital sinir bloğu.

Introduction

Ruptured aneurysms account for approximately 80% of non-traumatic SAH. The 30-day mortality rate of aneurysmal SAH is 45% and 30% of the survivors have significant neurological problems. SAH is classically manifested with headaches.

97% of the diagnosed patients have headache.^[1] Pain

can be observed in 25–56% of the ruptured or unruptured aneurysms due to EVT.^[2,3] It has been reported that causes of pain include; stretching and distortion of the arteries with coiling and stenting, and inflammatory changes caused by chemicals such as glue.^[4] It has also been shown that bioactive coils and the absence of hypertension history increase the risk of headache.^[3,5] Despite the use of agents such as ac-

Department of Anesthesiology and Reanimation, Başkent University Faculty of Medicine, Ankara, Turkey

Submitted: 30.03.2018 Accepted after revision: 05.09.2018 Available online date: 01.10.2018

Correspondence: Dr. Rafi Doğan, Başkent Üniversitesi Tıp Fakültesi, Anesteziyoloji ve Reanimasyon Anabilim Dalı, Ankara, Turkey.

Phone: +90 - 332 - 257 06 06 e-mail: rafidogan@yahoo.com

© 2018 Turkish Society of Algology

etaminophen, opioids, tramadol, and gabapentin for the treatment of headache; effective pain control has not been achieved and undesirable side effects of drugs have come to the forefront. These medicines may also prevent objective neurological evaluation by alteration of consciousness at different degrees.

Occipital nerve block has been shown to provide effective pain control in occipital neuralgia, cervicogenic headache, tension-type headache and migraine.^[6-9] US-guided GON block increases the efficiency and reliability of the block.^[10]

In this report we will present bilateral greater occipital nerve (GON) block that is performed to two patients who underwent EVT and stenting for a ruptured and unruptured intracranial aneurysm.

Case Report

Case 1 – A 59-year-old male patient with a history of hypertension was admitted to the hospital due to unconsciousness, computerized brain tomography showed an extensive SAH (Fisher 3), which also was opened to the ventricle. Glasgow coma score (GCS) was 13, other physical examinations were normal. Cerebral angiography showed aneurysm in anterior communicating artery. Lumbar spinal drainage catheter was inserted into the patient. After EVT, daily cerebrospinal fluid (CSF) drainage (30–40 ml/day) was performed.

Coilization and stenting were done with EVT. During the procedure, 2000 U of fractionated heparin was administered to the patient. After the procedure, the patient was admitted to the intensive care unit (ICU). After the patient gained consciousness the pain score (VAS) was assessed as 9–10 and we decided to perform bilateral GON block. When the patient was in prone position, the external occipital protuberance (EOP) was palpated, and then the 5–12 Hz linear US probe was placed in the transverse position on the midline EOP and advanced to the caudal. When the C2 vertebra was seen, the probe was shifted laterally and the oblique capitis inferior muscle (OCI) observed when the outer edge of the probe reached to the C1 transverse process level. GON was seen between OCI and semispinal capitis muscles and medial to occipital artery. The 25 G spinal needle was advanced in-plane from the medial to the lateral

towards the two muscle fascia. A mixture of 1 ml of 2% lidocaine, 2.5 ml of 0.25% bupivacaine and 3 mg of betamethasone (total 4 ml) was injected into the nerve sheath. Spread between two muscles was also observed with US.

After the block the pain score was 1. The VAS was questioned 30 minutes after the block and then every 12 hours for 14 days until discharge. Oral nimodipine 360 mg/day for vasospasm prophylaxis, and 5 mg prasugrel and 100 mg ASA for antiaggregant therapy were initiated. During these 14 days, if the pain score was 4 and above, paracetamol was administered 500 mg iv 4 times a day, and tramadol 100 mg iv was administered if the pain score did decrease. Control angiography was performed at 8th and 10th days because the patient became unconscious and experienced regression in motor functions. Cerebral vasospasm was observed and treated with intra arterial papaverine. After 14 days the patient was transferred to the neurology ward with minimal motor weakness on the right lower limb.

Case 2 – A 55-year-old male patient presented with left leg numbness and headache. The patient had hypertension and diabetes mellitus history. The headache was concentrated in the back of the neck, and described as constant and throbbing. GCS was 15, minimal sensory loss in left leg was observed and other systemic examinations were normal. Computerized brain tomography revealed a basilar artery aneurysm and small infarct in the brain stem but no SAH was observed.

Under general anesthesia, left basilar artery stenting and fusiform aneurysm coiling were performed with EVT. During the procedure, 2000 U of fractionated heparin was administered to the patient. The patient was taken to ICU after recovery from anesthesia. Because VAS score, obtained in the pain questionnaire, was 7; the patient was scheduled for bilateral US-guided GON block. And performed as described in first case. After the block, the patient's VAS score was 2. VAS did not exceed 3 during the follow-up in ICU, so there was no need for additional analgesics. The sensory deficit of the patient resolved. Oral nimodipine 360 mg/day for the vasospasm prophylaxis and 5 mg prasugrel and 100 mg ASA for antiaggregant therapy were initiated. The patient did not develop

vasospasm and was transferred to the neurology ward on the 12nd day.

Discussion

In the presented 2 cases, effective pain control was obtained with GON block. Although Case 1 required minimal additional analgesia, it can be said that the block was effective.

There are 3 reasons for headache in aneurysmatic SAH which are treated by EVT. The first reason is SAH itself, the second is EVT, and the last is the drainage of the BOS with lumbar drainage in cases of extensive SAH which are opened to the ventricle. These factors can cause pain, either alone or together. In other words, it is not possible to determine which factor is responsible at any given time. The cause of headache in SAH is not fully explained, but sudden stretching of the aneurysm, local meningeal inflammation or thrombosis is implicated.^[5,11] Pain is described as sudden, explosive and spreading to the neck. It usually takes 1–2 weeks and the pain control can be performed with only potent analgesics such as opioids and tramadol.

Sometimes very severe pain may also be resistant to continuous opioid therapy.^[11] In addition, the opioids can mask the neurological situation and the opioid antagonists are unable to remove this masking; which led researchers to search for different methods of pain control.^[12,13]

In SAH cases, the activation of NMDA receptors facilitate the transmission of pain leading to hyperalgesia.^[14] By considering this fact the analgesic efficacy of magnesium, an NMDA receptor antagonist, was investigated and shown that headache was less in patients with high Mg levels.^[15] There are studies showing that Mg replacement therapy is effective in some headache treatments.^[16] However, information on efficacy and safety of intravenous Mg support for pain control is limited. In addition, Mg was used as neuroprotective and antivasospastic in SAH patients. However, additional Mg supplementation is not recommended in this regard except for keeping serum levels within normal limits. In our study, serum Mg levels in both patients were within normal limits.

Although aneurysmatic SAH patients treated with

EVT are reported to have less headache,^[17] some studies reported headache related to EVT.^[2,4,5] In these patients it has been shown that causes of pain may be distortion, stretching of the vessel with coil or stent; inflammatory changes caused by chemicals such as glue.^[4] It has also been shown that bioactive coils and the absence of hypertension history increase the risk of headache.^[2,5] Evidence for the safety of tryptophan and dihydroergotamine, serotonin receptor antagonists is insufficient this type of headache.^[4]

Cerebral vasospasm and cerebral ischemia are the most serious complications after SAH, with a rate of 30–70%.^[18] It develops most frequently between 4–14 days.^[19] It is thought that hemoglobin products in CSF are effective in the development of vasospasm.^[20,21] The effectiveness of CSF drainage to prevent vasospasm has been demonstrated by many researchers. Lumbar drainage has been shown to be more efficient than external ventricular drainage for cleaning hemoglobin products from CSF.^[22,23] Although lumbar drainage can prevent vasospasm, it can also cause headache due to decreased CSF volume at the same time.

In our clinic, almost all of the ruptured or unruptured ICAs are treated with EVT. The procedure is performed under general anesthesia and the patient is followed up in the ICU. The duration of ICU stay is 10–15 days, especially due to the risk of vasospasm. Patients in ICU are also followed for headache treatment. According to the severity of headache, we use paracetamol, pregabalin, tramadol and opioids alone or in combination. We have observed many times that opioids are inadequate for pain control in severe headaches. Therefore, to increase efficacy in headache control and to move away from pregabalin, tramadol and opioids which have many side effects we decided to perform nerve blocks. We think that the most appropriate method for this purpose is the bilateral GON block.

GON block has been shown to provide effective pain control in occipital neuralgia, cervicogenic headache, tension-type headache and migraine.^[6–9] Performing the nerve block with the US guidance has increased the safety and effectiveness of the block. In a study in which the GON block was used

for the treatment of occipital neuralgia and cranio-facial pain syndromes, the efficacy of the block for 1 week, 1 month and 3 months was assessed and at least a 50% reduction in the VAS score was observed.^[10] They also emphasized that US is a safe and easily applicable imaging method. In another study, a significant reduction in pain scores over 4 weeks with GON block for cervicogenic headache and occipital neuralgia was observed and no adverse events were reported.^[24]

Furthermore, the efficacy of the block also has been shown in the post-dural puncture headache.^[25,26] We have emphasized in our study that one of the causes of the headache might be lumbar spinal drainage. In this case, we think that headache is caused by a similar mechanism with post-dural puncture headache.

So, which mechanisms have been effective in pain control with GON block? CSF volume changing caused by dural stretch in SAH may activate the trigeminal nucleus caudalis, causing increased activity in the trigeminal and greater occipital nerves.^[27] The other theories about the genesis of pain associated with intracranial endovascular procedures include mechanical stimulation of the arterial wall by traction, stretching, and distortion from materials used (balloons, catheters, coils, glue, stent) with subsequent triggering of the trigeminovascular pathway from trigeminal C fiber stimulation.^[28,29] Additionally, GON's relation to second-order neurons of the trigemino-cervical complex, which receive afferent inputs from trigeminal peripheral nerves of the face, offers insight into how GON block may contribute to relieving headaches that are more frontal than occipital in distribution.^[30,31] The analgesia obtained after GON block may be explained by the central neuromodulatory effect that causes decreased central sensitization as a result of the temporary interruption of afferent input to the dorsal roots and trigeminal nucleus.^[32]

Also in a retrospective review, adult patients with post-traumatic headaches had significant improvement in their headaches up to six months after GON block with xylocaine and methylprednisolone.^[33] Additionally a study showed GON block may have a role in management of spontaneous intracranial hypotension headache.^[27]

In conclusion, an efficient headache control was achieved with the bilateral GON block and the need for additional analgesic medication was severely reduced.

Conflict-of-interest issues regarding the authorship or article: None declared.

Peer-review: Externally peer-reviewed.

References

1. Abraham MK, Chang WW. Subarachnoid Hemorrhage. *Emerg Med Clin North Am.* 2016 Nov;34(4):901-916.
2. Choi KS, Lee JH, Yi HJ, Chun HJ, Lee YJ, Kim DW. Incidence and risk factors of postoperative headache after endovascular coil embolization of unruptured intracranial aneurysms. *Acta Neurochir (Wien).* 2014 Jul;156(7):1281-7.
3. Hwang G, Jeong EA, Sohn JH, Park H, Bang JS, Jin SC, et al. The characteristics and risk factors of headache development after the coil embolization of an unruptured aneurysm. *AJNR Am J Neuroradiol.* 2012 Oct;33(9):1676-8.
4. Baron EP, Moskowitz SI, Tepper SJ, Gupta R, Novak E, Husain MS, et al. Headache following intracranial neuroendovascular procedures. *Headache.* 2012 May;52(5):739-48.
5. Takigawa T, Matsumaru Y, Nakai Y, Nakamura K, Hayakawa M, Tsuruta W, et al. Bioactive coils cause headache and fever after endovascular treatment of intracranial aneurysms. *Headache.* 2012 Feb;52(2):312-21.
6. Aubry S, Kastler B, Bier V, Hadjidekov V, Hussein HH, Fergane B. Evaluation of the effectiveness of CT-guided infiltration in the treatment of Arnold's neuralgia. *Neuroradiology.* 2009 Mar;51(3):163-8.
7. Vincent MB, Luna RA, Scanduzzi D, Novis SA. Greater occipital nerve blockade in cervicogenic headache. *Arq Neuropsiquiatr.* 1998 Dec;56(4):720-5.
8. Young W, Cook B, Malik S, Shaw J, Oshinsky M. The first 5 minutes after greater occipital nerve block. *Headache.* 2008 Jul;48(7):1126-8.
9. Saadah HA, Taylor FB. Sustained headache syndrome associated with tender occipital nerve zones. *Headache.* 1987 Apr;27(4):201-5.
10. Zipfel J, Kastler A, Tatu L, Behr J, Kechidi R, Kastler B. Ultrasound Guided Intermediate Site Greater Occipital Nerve Infiltration: A Technical Feasibility Study. *Pain Physician.* 2016 Sep-Oct;19(7):E1027-34.
11. Morad AH, Tamargo RJ, Gottschalk A. The Longitudinal Course of Pain and Analgesic Therapy Following Aneurysmal Subarachnoid Hemorrhage: A Cohort Study. *Headache.* 2016 Nov;56(10):1617-1625.
12. Baskin DS, Hosobuchi Y. Naloxone reversal of ischemic neurological deficits in man. *Lancet.* 1981 Aug 8;2(8241):272-5.
13. Bell BA, Miller JD, Neto NG, O'Neill P, Laughton LM. Effect of naloxone on deficits after aneurysmal subarachnoid hemorrhage. *Neurosurgery.* 1985 Apr;16(4):498-501.

14. Woolf CJ, Thompson SW. The induction and maintenance of central sensitization is dependent on N-methyl-D-aspartic acid receptor activation; implications for the treatment of post-injury pain hypersensitivity states. *Pain*. 1991 Mar;44(3):293-9.
15. Dorhout Mees SM, Bertens D, van der Worp HB, Rinkel GJ, van den Bergh WM. Magnesium and headache after aneurysmal subarachnoid haemorrhage. *J Neurol Neurosurg Psychiatry*. 2010 May;81(5):490-3.
16. Mauskop A, Altura BT, Cracco RQ, Altura BM. Intravenous magnesium sulfate rapidly alleviates headaches of various types. *Headache*. 1996 Mar;36(3):154-60.
17. Hong CK, Joo JY, Kim YB, Shim YS, Lim YC, Shin YS, et al. The Course of Headache in Patients With Moderate-to-Severe Headache Due to Aneurysmal Subarachnoid Hemorrhage: A Retrospective Cross-Sectional Study. *Headache*. 2015 Jul-Aug;55(7):992-9.
18. Green DM, Burns JD, DeFusco CM. ICU management of aneurysmal subarachnoid hemorrhage. *J Intensive Care Med*. 2013;Nov-Dec;28(6):341-54.
19. Loch Macdonald R. Management of cerebral vasospasm. *Neurosurg Rev*. 2006 Jul;29(3):179-93.
20. Claassen J, Bernardini GL, Kreiter K, Bates J, Du YE, Copeland D, et al. Effect of cisternal and ventricular blood on risk of delayed cerebral ischemia after subarachnoid hemorrhage: the Fisher scale revisited. *Stroke*. 2001 Sep;32(9):2012-20.
21. Munoz-Guillen NM, Leon-Lopez R, Tunez-Finana I, Cano-Sanchez A. Del vasoespasmio a la lesion cerebral precoz: una nueva frontera en la investigacion de la hemorragia subaracnoidea. *Neurologia*. 2013;28:309—16.
22. Klimo P Jr, Kestle JR, MacDonald JD, Schmidt RH. Marked reduction of cerebral vasospasm with lumbar drainage of cerebrospinal fluid after subarachnoid hemorrhage. *J Neurosurg*. 2004 Feb;100(2):215-24.
23. Maeda Y, Shirao S, Yoneda H, Ishihara H, Shinoyama M, Oka F, Sadahiro H, Ueda K, Sano Y, Kudomi S, Hayashi Y, Shigeeda T, Nakano K, Koizumi H, Nomura S, Fujii M, Nomura S, Suzuki M. Comparison of lumbar drainage and external ventricular drainage for clearance of subarachnoid clots after Guglielmi detachable coil embolization for aneurysmal subarachnoid hemorrhage. *Clin Neurol Neurosurg*. 2013 Jul;115(7):965-70.
24. Pingree MJ, Sole JS, O'Brien TG, Eldrige JS, Moeschler SM. Clinical Efficacy of an Ultrasound-Guided Greater Occipital Nerve Block at the Level of C2. *Reg Anesth Pain Med*. 2017 Jan/Feb;42(1):99-104.
25. Akyol F, Binici O, Kuyruklu Yildiz U, Karabakan G. Ultrasound-guided bilateral greater occipital nerve block for the treatment of post-dural puncture headache. *Pak J Med Sci*. 2015 Jan-Feb;31(1):111-5.
26. Uyar Türkyilmaz E, Eryilmaz NC, Güzey NA, Moraloğlu Ö. Bilateral greater occipital nerve block for treatment of post-dural puncture headache after caesarean operations. *Rev Bras Anestesiologia*. 2016 Sep-Oct;66(5):445-50.
27. Niraj G, Critchley P, Kodivalasa M, Dorgham M. Greater Occipital Nerve Treatment in the Management of Spontaneous Intracranial Hypotension Headache: A Case Report. *Headache*. 2017 Jun;57(6):952-955
28. Martins IP, Baeta E, Paiva T, Campos J, Gomes L. Headaches during intracranial endovascular procedures: A possible model of vascular headache. *Headache*. 1993;33:227-233.
29. Beekman R, Nijssen PC, van Rooij WJ, Wijnalda D. Migraine with aura after intracranial endovascular procedures. *Headache*. 2001;41:410-413.
30. Bartsch T, Goadsby PJ. Increased responses in trigemino-cervical nociceptive neurons to cervical input after stimulation of the dura mater. *Brain*. 2003;126:1801-1813.
31. Busch V, Jakob W, Juergens T, Schulte-Mattler W, Kaube H, May A. Functional connectivity between trigeminal and occipital nerves revealed by occipital nerve blockade and nociceptive blink reflexes. *Cephalalgia*. 2006;26:50-55.
32. Bartsch T, Goadsby PJ. Stimulation of the greater occipital nerve induces increased central excitability of dural afferent input. *Brain*. 2002;125:1496—509.
33. Gawel MJ, Rothbart PJ. Occipital nerve block in the management of headache and cervical pain. *Cephalalgia*. 1992;12:9-13.