

Author's Reply

To the Editor,

In our study (1) limitations, we had already shown that the number of cases was small, and there may have been selection bias. In addition, this was a retrospective rather than a randomized controlled prospective study. All these shortcomings limited the value of the article. Whether such method can be successfully completed depended mainly on the images provided by transthoracic echocardiography (TTE). Perhaps because of the lean physique of the southern Chinese people, we found that using TTE could achieve satisfactory imaging for guiding device closure of atrial septal defect for most of the cases in our study. Meanwhile, some other papers supported the idea on device closure of ASD guiding by complete TTE (2, 3). However, for a few complex cases with poor images by TTE, we still used transesophageal echocardiography as the guiding tool. It is important to emphasize that we do not want to claim that the transthoracic method can replace the transcatheter method. The transthoracic method can be used as an alternative for those patients who are unable or unwilling to be exposed to radiation.

 Qiang Chen,  Hua Cao,  Gui-Can Zhang,  Liang-Wan Chen,
 Heng Lu,  Ling-Li Yu

Department of Cardiovascular Surgery, Union Hospital, Fujian Medical University; Fuzhou-*China*

References

1. Chen Q, Cao H, Zhang GC, Chen LW, Lu H, Yu LL. Transcatheter device closure of atrial septal defects guided completely by transthoracic echocardiography: A single cardiac center experience with 152 cases. *Anatol J Cardiol* 2018; 20: 330-5.
2. Bartakian S, El-Said HG, Printz B, Moore JW. Prospective randomized trial of transthoracic echocardiography versus transesophageal echocardiography for assessment and guidance of transcatheter closure of atrial septal defects in children using the Amplatzer septal occluder. *JACC Cardiovasc Interv* 2013; 6: 974-80.
3. Azhar AS. Safety and efficacy of transthoracic versus transesophageal echocardiography in transcatheter closure of atrial septal defects. Reporting a single center experience from Saudi Arabia. *Saudi Med J* 2016; 37: 1196-205.

Address for Correspondence: Qiang Chen, MD,

Department of Cardiovascular Surgery,

Union Hospital,

Fujian Medical University,

Xinquan Road 29# 362000

Fuzhou-*China*

Phone: +861 379 937 62 16

E-mail: chenqiang2228@163.com

©Copyright 2018 by Turkish Society of Cardiology - Available online at www.anatoljcardiol.com