

atrial strain and strain rate in the choice of proper treatment for patients with atrial fibrillation.

Karolina Kupczynska
Department of Cardiology, Medical University of Lodz, Bieganski Hospital, Lodz-Poland

References

1. Kupczynska K, Michalski BW, Miskowicz D, Kasprzak JD, Wejner-Mik P, Wdowiak-Okrojek K, et al. Association between left atrial function assessed by speckle-tracking echocardiography and the presence of left atrial appendage thrombus in patients with atrial fibrillation. *Anatol J Cardiol* 2017; 18: 15-22.
2. Kusunose K, Yamada H, Nishio S, Tomita N, Hotchi J, Bando M, et al. Index-beat assessment of left ventricular systolic and diastolic function during atrial fibrillation using myocardial strain and strain rate. *J Am Soc Echocardiogr* 2012; 25: 953-9.

Address for Correspondence: Dr. Karolina Kupczynska,
 Department of Cardiology, Medical University of Lodz
 Bieganski Hospital, Kniaziewiczza 1/5 91-347 Lodz-Poland
 Phone: +48 42 251 62 16 Fax: +48 42 653 99 09
 E-mail: karolinakupczynska@gmail.com

Ganglionated plexi interactions with atrio-ventricular node and right vagus nerve

To the Editor,

We read with great interest the article titled "Vagal denervation in atrial fibrillation ablation: A comprehensive review" by Aksu et al. (1) published in *Anatol J Cardiol* 2017; 18: 142-8. It is an excellent narrative review about ganglionated plexi (GP), vagal denervation, and atrial fibrillation. We have some commentaries about GP interactions with atrio-ventricular node (AVN) and right vagus nerve.

There is a large interactive network among different GP; this network serves as an "integrated center" of the cardiac autonomic innervation (2). Vagus nerve exerts its influence on the AVN through the epicardial fat pads that are primarily located on the posterior wall of the left atrium (2). The inferior vena cava-left atrium fat pad (namely also right inferior GP) located around the coronary sinus mainly provides vagal innervations and selectively innervates the AVN in humans (2, 3). It was shown that high-frequency stimulation of the right anterior (or superior right atrial vagal GP) and left superior GP (or superior left atrial vagal GP) could also influence the AVN (3). In addition, the influence of the right anterior GP on the AVN appears to be more important than its influence on the left superior GP (2, 3).

A functional neural pathway between the right vagus nerve and the AVN was identified (2), and the integrity of the GP seems to represent a mandatory interconnected network (3). In this study by Xhaet et al. (3), the absence of any alteration in the

ventricular rate in response to high-frequency stimulation of the right vagus nerve after the ablation of GP suggests that the right vagus nerve is not directly connected to the AVN and that the integrity of the GP is required to produce vagal effects on the AVN (3). Probably, there is no direct pathway between both the right and left vagus nerves and the AVN.

The long-term influence of GP ablation on the electrophysiology of the AVN is not known. However, the incomplete GP ablation can increase the vulnerability of the atria to atrial fibrillation and denervation is likely transient (4, 5). In addition, GP ablation that led to parasympathetic denervation of the AVN could play a role in the high ventricular rate response of atrial tachycardia after atrial fibrillation ablation. Therefore, GP interaction with the AVN and right vagus nerve could provide new insights on that particular mechanism.

Figure 1 is very interesting from the anatomical point of view, but there are two GC without any GB. The alignment of the GP nomenclature is also obviously required.

In conclusion, there are GP interactions with the AVN and right vagus nerve with possible important consequences on vagal denervation in atrial fibrillation ablation. However, the role and influence of the GP on the complicated vagal innervation of the heart still needs to be clarified.

Mariana Floria^{*,}, Olivier Xhaet¹, Ileana Antohe^{*,**}**
^{*}Sf. Spiridon Emergency Hospital, ^{**}Grigore T. Popa University of Medicine and Pharmacy, Iasi-Romania
¹Université catholique de Louvain, CHU UCL Namur-Belgium

References

1. Aksu T, Güler TE, Mutluer FO, Oto MA. Vagal denervation in atrial fibrillation ablation: A comprehensive review. *Anatol J Cardiol* 2017; 18: 142-8. [\[CrossRef\]](#)
2. Hou Y, Scherlag BJ, Lin J, Zhang Y, Lu Z, Truong K, et al. Ganglionated plexi modulate extrinsic cardiac autonomic nerve input: effects on sinus rate, atrioventricular conduction, refractoriness, and inducibility of atrial fibrillation. *J Am Coll Cardiol* 2007; 50: 61-8.
3. Xhaet O, DE Roy L, Floria M, Deceuninck O, Blommaert D, Dormal F, et al. Integrity of the Ganglionated Plexi Is Essential to Parasympathetic Innervation of the Atrioventricular Node by the Right Vagus Nerve. *J Cardiovasc Electrophysiol* 2017; 28: 432-7. [\[CrossRef\]](#)
4. Hirose M, Leatmanorath Z, Laurita KR, Carlson MD. Partial vagal denervation increases vulnerability to vagally induced atrial fibrillation. *J Cardiovasc Electrophysiol* 2002; 13: 1272-9. [\[CrossRef\]](#)
5. Oh S, Zhang Y, Bibevski S, Marrouche NF, Natale A, Mazgalev TN. Vagal denervation and atrial fibrillation inducibility: Epicardial fat pad ablation does not have long-term effects. *Heart Rhythm* 2006; 3: 701-8. [\[CrossRef\]](#)

Address for Correspondence: Mariana Floria, MD, PhD, FESC
 From IIIrd Medical Clinic and Grigore
 T.Popa University of Medicine and Pharmacy
 16 University Street, Iasi-Romania
 Phone: +40.232.301.600 Fax: +40.232.211.820
 E-mail: floria_mariana@yahoo.com

©Copyright 2017 by Turkish Society of Cardiology - Available online
 at www.anatoljcardiol.com
 DOI:10.14744/AnatolJCardiol.2017.8087

