

Myocardial infarction as a thrombotic complication of myeloproliferative disorders

Essential thrombocythemia (ET) and polycythemia vera (PV) are myeloproliferative disorders (MPDs), comprising a heterogeneous group of hematological diseases characterized by excessive production of cells belonging to the myeloid lineage in the bone marrow. They are related to, and may evolve into, myelodysplastic syndromes and acute myeloid leukemia, although MPDs as a whole have a much better prognosis than these conditions (1).

MPDs are typically complicated by thrombosis, with a rate of major events as high as 50% (2). Advanced age and a prior history of thrombosis are the two most important risk factors for vascular complications, although hypercholesterolemia, hypertension, smoking, and diabetes have been recognized as incremental factors of thrombosis in patients presenting with MPDs (3). More recently, leukocytosis has been reported as an independent risk factor for thrombosis in both ET and PV (4). Furthermore, the additional value of the JAK2 V617F mutation as a risk factor for thrombosis is currently being investigated (5).

The mechanisms behind the increased thrombotic tendency in MPDs are complex and remain to be elucidated. No correlation has been found between platelet values and the risk of thrombosis (6). Pathophysiological mechanisms possibly involve all cellular blood components, plasma, interaction with endothelial cells, as well as hemodynamic alterations. In MPDs, platelets exhibit the feature of spontaneous aggregation (7), and, in addition, they demonstrate abnormal expression of glycoprotein (8). Leukocytes also contribute to platelet activation in MPDs (9), and erythrocytes show an increased tendency to adhere to the endothelium (10). Interestingly, leukocyte count seems to be a stronger predictor for thrombosis than do platelet counts and hemoglobin/hematocrit levels (9). The hemodynamic changes include increased viscosity, with displacement of circulating platelets toward the endothelium (11). Increased levels of microparticles with procoagulant activity are also found in MPD patients (12). Increased cytokine expression, leading to a general inflammatory response, probably also contributes to the thrombotic tendency (13). An increased number of prothrombotic circulating endothelial cells are seen among these patients; notably, the JAK2 V617F mutation is found in these cells (14).

In these respects, the paper entitled "Myocardial infarction as a thrombotic complication of essential thrombocythemia and polycythemia vera" by Pósfai et al. (15) published in the cur-

rent issue of the Anatolian Journal of Cardiology is of potential interest. In this descriptive study, 263 patients diagnosed with either ET or PV were included. In a retrospective analysis, the clinical characteristics of 14 patients who suffered myocardial infarction (MI) during their hematological follow-up were compared to 162 MPD patients who did not exhibit any kind of thrombotic complication. A major result of the study is that most of the MI complications appeared within 1 year after the hematological diagnosis of ET or PV, with evident implications for the necessity of early detection and management in this group of patients. Furthermore, JAK2 V617F mutation was present in most of the cases, and cardiovascular risk factors appeared in the majority of patients. Other interesting results refer to the variables of mean peripheral platelet count, hemoglobin, hematocrit, and red blood cell count.

Emmanouil Petrou, Vasiliki Karali¹

Division of Cardiology, Onassis Cardiac Surgery Center, Athens-Greece

¹First Department of Propaedeutic and Internal Medicine, Athens University Medical School, Athens-Greece

References

1. Tefferi A, Thiele J, Orazi A, Kvasnicka HM, Barbui T, Hanson CA, et al. Proposals and rationale for revision of the World Health Organization diagnostic criteria for polycythemia vera, essential thrombocythemia, and primary myelofibrosis: recommendations from an ad hoc international expert panel. *Blood* 2007; 110: 1092-7.
2. Elliott MA, Tefferi A. Thrombosis and haemorrhage in polycythemia vera and essential thrombocythaemia. *Br J Haematol* 2005; 128: 275-90. [[Crossref](#)]
3. Marchioli R, Finazzi G, Landolfi R, Kutti J, Gisslinger H, Patrono C, et al. Vascular and neoplastic risk in a large cohort of patients with polycythemia vera. *J Clin Oncol* 2005; 23: 2224-32. [[Crossref](#)]
4. Landolfi R, Di Gennaro L, Barbui T, De Stefano V, Finazzi G, Marfisi R, et al. Leukocytosis as a major thrombotic risk factor in patients with polycythemia vera. *Blood* 2007; 109: 2446-52. [[Crossref](#)]
5. Finazzi G, Rambaldi A, Guerini V, Carobbo A, Barbui T. Risk of thrombosis in patients with essential thrombocythemia and polycythemia vera according to JAK2 V617F mutation status. *Haematologica* 2007; 92: 135-6. [[Crossref](#)]
6. Besses C, Cervantes F, Pereira A, Florensa L, Solé F, Hernández-Boluda JC, et al. Major vascular complications in essential thrombocythaemia: a study of the predictive factors in a series of 148 patients. *Leukemia* 1999; 13: 150-4. [[Crossref](#)]

Address for correspondence: Emmanouil Petrou, MD, Division of Cardiology, Onassis Cardiac Surgery Center, 356 Syggrou Ave. Athens, GR-17476-Greece
E-mail: emmgpetrou@hotmail.com

Accepted Date: 17.03.2016 **Available Online Date:** 25.04.2016

©Copyright 2016 by Turkish Society of Cardiology - Available online at www.anatoljcardiol.com
DOI:10.14744/AnatolJCardiol.2016.18293



7. Ginsburg AD. Platelet function in patients with high platelet count. *Ann Intern Med* 1975; 82: 506-11. [\[Crossref\]](#)
8. Jensen MK, de Nully Brown P, Lund BV, Nielsen OJ, Hasselbalch HC. Increased platelet activation and abnormal membrane glycoprotein content and redistribution in myeloproliferative disorders. *Br J Haematol* 2000; 110: 116-24. [\[Crossref\]](#)
9. Barbui T, Carobbio A, Rambaldi A, Finazzi G. Perspectives on thrombosis in essential thrombocythemia and polycythemia vera: is leukocytosis a causative factor? *Blood* 2009; 114: 759-63. [\[Crossref\]](#)
10. Wautier MP, El Nemer W, Gane P, Rain JD, Cartron JP, Colin Y, et al. Increased adhesion to endothelial cells of erythrocytes from patients with polycythemia vera is mediated by laminin alpha5 chain and Lu/BCAM. *Blood* 2007; 110: 894-901. [\[Crossref\]](#)
11. Pearson TC, Wetherley-Mein G. Vascular occlusive episodes and venous haematocrit in primary proliferative polycythaemia. *Lancet* 1978; 2: 1219-22. [\[Crossref\]](#)
12. Duchemin J, Ugo V, Ianotto JC, Lecucq L, Mercier B, Abgrall JF. Increased circulating proagulant activity and thrombin generation in patients with myeloproliferative neoplasms. *Thromb Res* 2010; 126: 238-42. [\[Crossref\]](#)
13. Barbui T, Carobbio A, Finazzi G, Vannucchi AM, Barosi G, Antonioli E, et al. Inflammation and thrombosis in essential thrombocythemia and polycythemia vera: different role of C-reactive protein and pentraxin 3. *Haematologica* 2011; 96: 315-8. [\[Crossref\]](#)
14. Reikvam H, Tiu RV. Venous thromboembolism in patients with essential thrombocythemia and polycythemia vera. *Leukemia* 2012; 26: 563-71. [\[Crossref\]](#)
15. Pósfai E, Marton I, Borbényi Z, Nemes A. Myocardial infarction as a thrombotic complication of essential thrombocythaemia and polycythaemia vera. *Anatol J Cardiol* 2016; 16: 397-402.