

**Ahmet Korkmaz, Havva Tuğba Gürsoy, Bekir Demirtaş, Özgül Uçar Elalmış, Ümit Güray**  
**Department of Cardiology, Ankara Numune Training and Research Hospital; Ankara-Turkey**

## References

1. Robinson T, Oliver J, Sheridan P, Sahu J, Bowes R. Fragmentation and embolization of pacemaker leads as a complication of lead extraction. *Europace* 2010; 12: 754-5.
2. Kennergren C. A European perspective on lead extraction: Part I. *Heart Rhythm* 2008; 5: 160-2.
3. Smith MC, Love CJ. Extraction of transvenous pacing and ICD leads. *Pacing Clin Electrophysiol* 2008; 31: 736-52.
4. Byrd C, Wilkoff B, Love CJ, Sellers TD, Reiser C. Clinical study of the laser sheaths for lead extraction: the total experience in the United States. *Pacing Clin Electrophysiol* 2002; 25: 804-8.

**Address for Correspondence:** Dr. Ahmet Korkmaz,  
 Ankara Numune Eğitim ve Araştırma Hastanesi,  
 Kardiyoloji Bölümü, Altındağ, Ankara-Türkiye  
 Phone: +90 312 508 40 00  
 E-mail: drahmtrkmz07@gmail.com



©Copyright 2018 by Turkish Society of Cardiology - Available online  
 at [www.anatoljcardiol.com](http://www.anatoljcardiol.com)  
 DOI:10.14744/AnatolJCardiol.2017.8249

## Ten years' clinical experience of cardiac myxoma: diagnosis, treatment, and clinical outcomes

To the Editor,

Cardiac myxoma (CM) is the most common type of benign primary cardiac tumor (1). Approximately more than half of primary cardiac tumors are myxomas (2). They are most commonly diagnosed between the age of 30 and 60 years (3). CM is described as a sporadic or familial disorder in the literature (2, 3). A limited number of patients have been referred with the classical triad of obstructive cardiac symptoms: pulmonary edema, progressive heart failure (HF), and arterial embolic events. Rarely, syncope/vertigo or sudden death can be the first symptom of CM. For early diagnosis, transthoracic echocardiography (TTE) is being increasingly used. Recently, magnetic resonance imaging (MRI) and/or thoracic computed tomography (CT) have been used for prompt diagnosis. Early and optimal surgical excisions have shown excellent early- and long-term results, with no recurrence of the tumor (4). According to previous studies, CMs may be diagnosed sporadically in 90% of patients (5).

In contrast to solid myxoma, papillary myxoma is characterized by a soft formation that is friable during tumor excision. Therefore, the rate of tumor recurrence is high in patients with papillary myxoma than in those with solid myxoma (4).

We treated 38 patients with CMs between June 2006 and September 2016 and retrospectively analyzed the symptoms,

diagnostic methods, and treatment strategies. Briefly, the mean age of the patients who underwent primary myxoma resection was 41.7±7.8 years, and female/male ratio was 22/16. Two patients with CM were in the pediatric age group (13 and 17 years). We used two-dimensional TTE for the diagnosis of CM in all patients. If tumors other than myxomas were suspected, thoracic CT or MRI was used.

No mortality occurred in the early postoperative period. Three patients required an emergent operation because of HF. In the early postoperative period, we detected a low cardiac output syndrome, new onset of atrial fibrillation, and mediastinal bleeding in 12 patients. Mean ICU and length of hospital stay was 2.7±1.4 and 8.5±3.3 days, respectively. Two patients died at a mean follow-up of 32±13 months postoperatively. Among the 36 long-term survivors, 76% of patients were in NYHA class I, whereas 24% were in NYHA class II. Two patients who underwent left atrial myxoma resection showed a recurrence 33 and 46 months after the first surgery. Congestive HF resulting from obstructive cardiac manifestations was detected in seven patients.

CM can present with a wide range of symptomatic spectrum from being asymptomatic to having serious side effects (4, 5). Our patients with a large solid myxoma that were localized in the left atrium had a greater incidence of HF and obstructive symptoms. In accordance with our experiences, serious proteinuria or acute renal failure which may be the first sign of right atrial myxoma. Peripheral or cerebral artery embolic events are the main catastrophic symptoms related to tumor type and location. After diagnosis is confirmed, early tumor excision should be performed. Surgery has excellent overall survival and freedom from reoperation, but follow-up using TTE is recommended.

**Hidayet Kayañçiçek, Emjed Khalil<sup>1</sup>, Gökhan Keskin<sup>2</sup>, Özkan Alataş<sup>3</sup>, Erhan Hafız<sup>4</sup>, Ömer Faruk Doğan<sup>5</sup>**

**Department of Cardiology, Private Medical Park Elazığ Hospital; Elazığ-Turkey**

<sup>1</sup>The Health Sciences University, Ersin Aslan Research and Training Hospital; Gaziantep-Turkey

<sup>2</sup>Department of Cardiology, Amasya University Research and Training Hospital; Amasya-Turkey

<sup>3</sup>Department of Radiology, The Health Sciences University, Elazığ Research and Training Hospital; Elazığ-Turkey

<sup>4</sup>Department of Cardiovascular Surgery, The Health Sciences University, Ersin Aslan Research and Training Hospital; Gaziantep-Turkey

<sup>5</sup>Department of Cardiovascular Surgery, The Health Sciences University, Adana Numune Research and Training Hospital; Adana-Turkey

## References

1. Garatti A, Nano G, Canziani A, Gagliardotto P, Mossuto E, Frigiola A, et al. Surgical excision of cardiac myxomas: twenty years experience at a single institution. *Ann Thorac Surg* 2012; 93: 825-31.
2. Keeling IM, Oberwalder P, Anelli-Monti M, Schuchlenz H, Demel U, Titz GP, et al. Cardiac myxomas: 24 years of experience in 49 patients. *Eur J Cardiothorac Surg* 2002; 22: 971-7. [CrossRef]
3. Steger CM, Hager T, Ruttman E. Primary cardiac tumours: a single-

- center 41-year experience. *ISRN Cardiol* 2012; 2012: 906109.
4. Tansel T, Harmandar B, Ugurlucan M, Nisli K, Eker R, Sozen AB, et al. Over 14 years of experience with cardiac myxomas. *Acta Cardiol* 2006; 61: 285-8. [CrossRef]
  5. Zheng JJ, Geng XG, Wang HC, Yan Y, Wang HY. Clinical and histopathological analysis of 66 cases with cardiac myxoma. *Asian Pac J Cancer Prev* 2013; 14:1743-6. [CrossRef]

**Address for Correspondence:** Dr. Ömer Faruk Doğan,  
Adana Numune Eğitim ve Araştırma Hastanesi,  
Adana, 01647-Türkiye

Phone: +90 533 481 30 56

Fax: +90 322 355 01 01

E-mail: ofdogan@hacettepe.edu.tr

©Copyright 2018 by Turkish Society of Cardiology - Available online  
at [www.anatoljcardiol.com](http://www.anatoljcardiol.com)

DOI:10.14744/AnatolJCardiol.2017.8265



## Sudden cardiac death in physicians: an alarming problem in China

To the Editor,

The recent revised declaration of Geneva highlights that physicians should attend to their own health, well-being, and abilities in order to provide care of the highest standard (1). Nevertheless, in China, sudden cardiac death has frequently occurred in the past 2 years. In our statistical analysis, the number of physician deaths was 17 and 23 in 2016 and 2017, respectively, which were notably increased compared with the reported 29 deaths from November 2008 to April 2015 (2). The percentage of male physicians was 95.7%. The mean age was 36.30±8.41 years. In this year, sudden cardiac death occurred in nine surgeons, five anesthesiologists, six internal physicians, and three intervention specialists. A 24-h shift or a night shift (11 cases) and preparing a ranking examination (7 cases) were the most common causes.

Male physicians, especially physicians in surgery and operative care, have larger overloads and longer working hours, which may be the primary cause of death. Besides, incorrect stress relief manners also affect physicians' healths, such as excessive alcohol consumption, smoking, and long-term energy drink consumption (3). Lack of exercise and obesity also increase the risk of acute myocardial infarction and stroke. Physicians usually neglect regular medical checkup themselves, which results in some potential diseases not being timely detected, such as hypertension, hyperlipidemia, and hyperuricemia.

The incidence of sudden death was also closely associated with economic and social factors. Middle-aged physicians

have heavy familial responsibilities, such as supporting parents, rearing children, and mortgage payments, which cause them to work harder. Deteriorated working environment, serious violence against doctors, and decreased patient-physician trust also intensified the sense of "burnout" in Chinese doctors (4, 5). Besides, because of the uneven distribution of healthcare resources in China, more than 70% of deaths were of physicians from tertiary hospitals.

We suggest that physicians should take responsibility for their own health. They should regularly perform aerobic exercise or connect with families and friends for support, instead of excessive smoking and drinking. Besides, regular medical checkup is an effective approach to decrease the risk of some potential diseases such as coronary artery disease, hypertension, and stroke. Government and media should help to improve the working environment and re-establish patient-physician trust, which may further decrease the sense of "burnout" for physicians (4, 5). Imperative changes for the assessments of physicians in China are also required to decrease the pressure of publishing papers or frequently preparing professional examinations.

**Xinmiao Shi, Rui Liu<sup>1</sup>**

**Department of Pediatrics, Peking University First Hospital;  
Beijing-China**

**<sup>1</sup>Heart Center, The First Hospital of Tsinghua University;  
Beijing-China**

### References

1. Parsa-Parsi RW. The Revised Declaration of Geneva: A Modern-Day Physician's Pledge. *JAMA* 2017; 318: 1971-2.
2. Song XN, Shen J, Ling W, Ling HB, Huang YM, Zhu MH, et al. Sudden Deaths Among Chinese Physicians. *Chin Med J (Engl)* 2015; 128: 3251-3.
3. Kaşıkçıoğlu E. Sports, energy drinks, and sudden cardiac death: stimulant cardiac syndrome. *Anatol J Cardiol* 2017; 17:163-4.
4. Yang Y, Zhao JC, Zou YP, Yan LN. Facing up to the threat in China. *Lancet* 2010; 376: 1823.
5. Lyu Z, Wu S, Cai Z, Guan X. Patient-physician trust in China: health education for the public. *Lancet* 2016; 388: 2991.

#Xinmiao Shi and Rui Liu contribute equally to the article.

**Address for Correspondence:** Xinmiao Shi, MD, PhD,  
Department of Pediatrics, Peking University First Hospital,  
No.1 Xi An Men Da Jie, Beijing-China

Phone: +86 10 8357 3238

Fax: +86 10 6653 0532

E-mail: shixinmiaozhenhao@163.com

©Copyright 2018 by Turkish Society of Cardiology - Available online  
at [www.anatoljcardiol.com](http://www.anatoljcardiol.com)

DOI:10.14744/AnatolJCardiol.2017.8263

