

Single coronary artery diagnosed by multidetector computed tomography

Çok kesitli bilgisayarlı tomografi ile tanı konulan tek koroner arter

The coronary circulation originating from a single coronary ostium is a rare variation. These variations were classified according to the site of the origin and distribution patterns of the coronary arteries. We describe Type IIB 4 pattern in this case. The left main coronary artery (LMCA) arised from the proximal right coronary artery and coursed to the left side anterior to the right ventricle. It gave the left anterior descending artery (LAD) coursing to the interventricular groove and left circumflex artery (CxA) coursing to the left atrioventricular groove.

A 40-year-old male presented with atypical chest pain for six month. His physical examination, laboratory results, chest X-ray, and electrocardiogram were normal. A multidetector computed tomography coronary angiography (MDCT-CA) was performed to evaluate the coronary vessels. The MDCT-CA showed a single large coronary artery originating from right sinus of Valsalva (Fig. 1, Video 1. See corresponding video/images at www.anakarder.com) and the left main coronary artery arising from proximal right coronary artery. After traversing in front of the right ventricular outflow tract, LMCA gave the origin to the LAD and CxA.



Figure 1. View of a single coronary artery originating from right coronary cusp in 3D- volume rendered by multidetector computed tomography image



Figure 2. Absence of vessels originating from left coronary cusp in volume rendered by multidetector computed tomography image

There was no vessel originating from the left coronary cusp (Fig. 2). Stenosis was not observed in any part of the coronary tree. No further associated congenital heart anomaly was present.

The incidence of isolated single coronary artery in angiographic series is 0.024%. It has a little clinical significance, except for cases in which a coronary artery traverses between the aorta and the pulmonary artery. Identification of retropulmonary position of an arterial segment is important, since mechanical compression of such a vessel between the aorta and the pulmonary artery may be a potential cause of ischemia and sudden cardiac death. The anomaly of this case has no clinical significance of this kind.

Until now, the main diagnostic method for detection of coronary anomalies has been coronary angiography which has some limitations. It is an invasive procedure and only selectively visualizes one vessel tree at a time and cannot obtain information on the 3-dimensional (3D) course of the vessel. Novel advances on computed tomography technology offer a noninvasive and accurate diagnostic modality to visualize the origin and course of anomalous coronary arteries in a 3D fashion.

Cihan Duran*, Demet Erciyes, Mustafa Şirvancı*,
Murat Şener**, Murat Gülbaran*,***
From Departments of Radiology* and Cardiology**,
Florence Nightingale Hospital, ***Department of Cardiology,
Faculty of Medicine, İstanbul Bilim University, İstanbul, Turkey**

Address for Correspondence/Yazışma adresi: Dr. Cihan Duran
Radyoloji Departmanı Florence Nightingale Hastanesi
Abide-i Hürriyet Cad.No:290 Şişli/İstanbul 80220, Turkey
Fax: +90 212 224 49 50/5010 Mobile: +90 532 667 60 56
E-mail: cduran65@myynet.com

Coronary artery evaluation in Kawasaki disease by dual source multi-detector CT coronary angiography in children

Çocuk yaşta Kawasaki hastalarında “dual source” multi dedektör BT anjiyografi ile koroner arter değerlendirmesi

Kawasaki disease is a childhood vasculitis of medium-sized vessels, affecting the coronary arteries in particular. Assessment of the degree of coronary stenosis and serial evaluation of coronary artery pathologies are essential for risk stratification and therapeutic management. Invasive coronary angiography is considered to be the gold standard technique. Multi-detector computed tomography coronary angiography (CTCA) is a noninvasive alternative imaging modality. It can also provide complementary morphologic data to echocardiography.

The CTCA is an approved noninvasive technique in the diagnosis of coronary pathologies. Dual source CTCA is a new generation device that can acquire diagnostic images in patients with high heart rates and/or arrhythmias without the use of beta blockers. This is advantageous for pediatric patients whose heart rates are generally higher than the average adult groups.

A 4-year old boy with 7 months' history of Kawasaki disease was admitted to the hospital for CTCA. In the previous echocardiographic examination 2 coronary aneurysms were detected in left main coronary artery (LMCA) and right coronary artery (RCA).

The CTCA was performed with a dual source (2x64 detector system) CT scanner during free breathing and injection of 40 mL of intravenous

contrast material. The average heart rate during the scan was 95 beats/min. No beta blocker was administered. Multiplanar reconstruction images showed three aneurysms; one in the distal LMCA, one in the mid-distal left anterior descending artery and the other one in the mid RCA (Fig. 1-4, Video 1. See corresponding video/movie images at www.anakarder.com). There was a striking fibrointimal proliferation in the RCA aneurysm causing luminal stenosis (Fig. 1, 2). No pericardial, myocardial or endocardial pathology was detected.




Figure 1. A volume rendered (VRT) axial 3D image showing the aneurysm in cross section and luminal stenosis with fibrointimal proliferation inner side (orange color) leaving a small lumen (white color)




Figure 2. A long-axis multiplanar image revealing the RCA aneurysm together with the intimal proliferation inner side

RCA- right coronary artery




Figure 3. A 3D volume rendered image showing the LMCA, LAD and CX aneurysms

LAD- left anterior descending artery, LMCA- left main coronary artery, CX- left circumflex artery




Figure 4. A 3D volume rendered image showing the RCA aneurysm

RCA- right coronary artery

Serkan Gelmez, Arda Saygılı*, Ercan Tutar**, Muzaffer Olcay Çizmeli
From Departments of Radiology and *Pediatric Cardiology,
Acıbadem Hospital, İstanbul
**Department of Pediatric Cardiology, Faculty of Medicine,
Ankara University, Ankara, Turkey

Address for Correspondence/Yazışma Adresi: Dr. Serkan Gelmez
Tekin Sok. No 111 Kadıköy İstanbul, Turkey
Phone: +90 216 544 43 44 Mobile: + 90 542 288 40 27 E-mail: sergelmez@yahoo.com

İlginç bir koroner arter fistülü olgusu: Diyagonal arter, sirkumfleks arter ve sağ koroner arter ile sol ventrikül arasında fistüller

An interesting case of coronary artery fistula: diagonal artery-, left circumflex artery-and right coronary artery to left ventricle fistulas

İlk kez 1865 yılında Krauze tarafından tanımlanan ve sıklığı %0.13 olarak bildirilen koroner arter fistülleri, koroner arterler ile kalp boşlukları, büyük damarlar (pulmoner arter, bronşiyal arter) veya diğer vasküler yapılar (mediyastinal damarlar, koroner sinüs) arasındaki anormal bağlantılardır.

Koroner arter fistüllerinin yaklaşık %50'sinin sağ koroner arterden, %42'sinin sol koroner arterden, %5'inin her iki koroner arterden kaynaklandığı, en sık sağ ventriküle (%41), sağ atriyuma (%26) ve pulmoner artere (%17), seyrek olarak da koroner sinüse (%7), sol atriyuma (%5) ve sol ventriküle (%3) açıldığı bildirilmiştir. Yaptığımız literatür taramasında aynı hastada üç koroner arterden sol ventriküle fistül hiç bildirilmemiştir.

Hipertansiyon ve diyabetes mellitus nedeni ile takipte olan 76 yaşında bayan hasta, stabil angina pectoris nedeniyle kliniğimize başvurdu. Fizik muayenede arteriyel kan basıncı 130/80 mm Hg, nabız sayısı 72 atım/dakika, düzenli idi. Apekte ve sol 3.-4. interkostal aralıkta 2/6 şiddetinde devamlı üfürüm tespit edildi. Diğer sistem muayenelerinde anormal bir bulguya rastlanmadı. Laboratuvar değerlerinde, telekardi-yografide ve ekokardiyografide anormal bulgu saptanmadı. Elektrokardiyografide sinüs ritmi, V 4-6 derivasyonlarında ST depresyonu izlendi. Efor testi pozitif bulunması üzerine yapılan koroner anjiyografide diyagonal