

Cardiogenic shock due to occlusion of left main coronary in a cocaine user

To the Editor,

Cocaine use has been related to the occurrence of myocardial infarction in young patients without other coronary risk factors. Acute myocardial infarction (AMI) secondary to the occlusion of the left main coronary artery (LMCA) in a cocaine user is infrequent, with sudden death being the most common form of presentation.

We present the clinical case of a 38-year-old male patient with an ST-segment elevation myocardial infarction complicated by cardiogenic shock because of acute occlusion of the LMCA following cocaine abuse.

He had a history of smoking and had recently used cocaine. He visited the emergency department of another institution with angina lasting for 5 h. The electrocardiogram (ECG) showed a heart rate of 120 bpm, complete RBBB with ST-segment elevation in lead aVR, lead I, aVL, and V2 to V6. The patient evolved with cardiogenic shock requiring mechanical ventilation (MV) and inotropic support. Considering the diagnosis of STEMI complicated with cardiogenic shock, the patient underwent coronary angiography. An intra-aortic balloon pump was placed before the procedure. The coronary angiography demonstrated a total acute thrombotic occlusion of the LMCA, and PPCI was performed. After the predilatation of the total occlusion with a balloon, TIMI-III flow was restored. A stent was successfully implanted.

After the procedure, the patient developed multiorgan failure (acute renal failure, liver failure, respiratory distress). Twenty-four hours later, he presented with ventricular tachycardia–ventricular fibrillation refractory, and the patient died.

In the present case, cocaine was presumed to be instrumental in provoking the AMI (patient with AMI was younger without classic risk factors). Cocaine stimulates the sympathetic nervous system by inhibiting catecholamine reuptake at sympathetic nerve terminals. Among them, include AMI and where the etiology is multifactorial (vasospasm, coronary dissection, atherosclerosis-plaque rupture, increased the determinants of myocardial oxygen consumption) (1).

Secondly, an acute obstruction of the LMCA is encountered at angiography approximately in only 0.5% of AMI cases (2), and it is associated with cardiogenic shock (2-4) as well as sudden death (5).

In patients with cardiogenic shock at admission, mortality was up to 32%–54% (3, 4). In patients with cardiogenic shock and multiorgan failure, mortality was up to 75% (4).

Coronary artery bypass graft surgery (CABG) is the standard revascularization strategy. However, normal blood flow in the infarct-related artery should be restored as rapidly and completely as possible; the high rate of mortality and of postoperative complications in patients with cardiogenic shock makes primary coronary intervention an alternative therapy. Percutaneous coronary intervention allows a rapid reperfusion of the vessel with a survival rate of 89% at 1 year (3, 4).

Among the variables associated with adverse outcomes, our patient presented with cardiogenic shock and underwent reperfusion therapy after 12 hours of symptom onset and multiorgan failure.

In conclusion, the etiology of AMI in patients with cocaine use is multifactorial. The occlusion of the LMCA is associated with high mortality secondary to cardiogenic shock. Survival depends on early reperfusion, and the appropriate strategy should be chosen based on the patient's hemodynamic status.

Gabriel E. Pérez Bazarrica, Mario L. Santa Cruz, Juan P. Arellano, Rafael Porcile
Department of Cardiology and Physiology, University Hospital, Faculty of Medicine, Universidad Abierta Interamericana; Buenos Aires-Argentina

References

1. Schwartz B, Rezkalla S, Kloner RA. Cardiovascular effects of cocaine. *Circulation* 2010; 122: 2558-69. [\[CrossRef\]](#)
2. Aygül N, Salamov E, Doğan U, Tokaç M. Acute occlusion of the left main trunk presenting as ST-elevation acute coronary syndrome. *J Electrocardiol* 2010; 43: 76-8. [\[CrossRef\]](#)
3. Pedrazzini G, Radovanovic D, Vassalli G, Sürder D, Moccetti T, Eberli F, et al. Primary percutaneous coronary intervention for unprotected left main disease in patients with acute ST-segment elevation myocardial infarction. *JACC Cardiovasc Interv* 2011; 6: 627-33. [\[CrossRef\]](#)
4. Pappalardo A, Mamas M, Imola F, Ramazzotti V, Manzoli A, Prati F, et al. Percutaneous coronary intervention of unprotected left main coronary artery disease as culprit lesion in patients with acute myocardial infarction. *JACC Cardiovasc Interv* 2011; 6: 618-6. [\[CrossRef\]](#)
5. Fiol M, Carrillo A, Rodriguez A, Pascual M, Bethencourt A, Bayés de Luna A. Electrocardiographic changes of ST-elevation myocardial infarction in patients with complete occlusion of the left main trunk without collateral circulation: differential diagnosis and clinical considerations. *J Electrocardiol* 2012; 45: 487-90. [\[CrossRef\]](#)

Address for Correspondence: Gabriel Pérez Bazarrica MD

Portela 2975 (1437). Buenos Aires-Argentina

Phone: (054)1149187561

E-mail: gpbazarrica@yahoo.com

©Copyright 2015 by Turkish Society of Cardiology - Available online

at www.anatoljcardiol.com

DOI:10.5152/AnatolJCardiol.2015.6409



An alternative malpractice system suggestion for Turkey: Patient compensation system

It is dangerous to be right in matters where established men are wrong

~ Voltaire ~

To the Editor,

Physicians and patients have started to realize that Turkish medical laws that enforced high medical malpractice compensation fines and sentenced physicians to imprisonment because of unintentional negligence are ruining the medical profession and healthcare system. If the present system continues on this track, physician burn out, increasing practice of defensive medicine, increasing cost of healthcare, and increasing mortality rates will be seen. In a widely referenced report, the cost of defensive medicine in USA is estimated to be USD 55.6 billion, which is equivalent to 2.4% of the health expenditure in 2008 (1). Unnecessary diagnostic tests and consultations and avoidance of high-risk patients are the most common form of defensive medicine (2). We have limited studies but some signs warn us that Turkey will face same consequences due to medical malpractice laws as long-lasting USA experience shows. It is needless to go through the same processes as USA for an additional 10–20 years in Turkey and face similar studies, discussions, high healthcare costs, and patient damages due to defensive medicine. We propose a new “patient compensation system” (PCS) for Turkey to avoid going through the same exhausting 20 years in the future.