

# What is cerebral palsy - new advances in treatment of cerebral palsy?

Milivoj Veličković Perat

*Academy of Developmental Medicine, Slovenia*

**Abstract.** Cerebral palsy is the most frequent motor impairment in childhood. In many countries, its prevalence is predicted to increase. There are an estimated 15 million people with cerebral palsy around the world and more than half of them are mentally retarded and one third have epilepsy. It should be seen that it is more than merely a motor disorder. There can be problems of development, perceptual-cognitive impairment, social and functional problems of daily living, emotional and behavioural problems, and many other difficulties. In addition to this early brain damage is not a synonym for Cerebral palsy.

Key words: Cerebral palsy, motor disorder, epilepsy, mental retardation

## 1. Introduction

Cerebral palsy (CP) is a well-recognized neurodevelopment condition beginning in early childhood and persisting through the lifespan. It was originally reported by J. Little in 1861 (but not then called 'cerebral palsy'), CP has been the subject of books and papers by some of the most eminent medical minds of the past hundred years. Beginning at the end of the 19th century, Sigmund Freud and Sir William Osler both contributed important perspectives on the condition. From the mid-1940s, the founding fathers of the American Academy for Cerebral Palsy and Developmental Medicine (Carlson, Crothers, Deaver, Fay, Perlstein, and Phelps) in the United States, and Mac Keith, Polani, Bax and Ingram of the Little Club in the United Kingdom, were among the leaders who moved the concepts and descriptions of CP forward.

### 1. 1. Definition of cerebral palsy

It has always been a challenge to define 'cerebral palsy'. One of definition used at present is definition of Cerebral Palsy from the Washington Workshop 2004: "Cerebral palsy describes a group of developmental disorders of movement and posture, causing activity restriction or disability, that are

attributed to disturbances occurring in the fetal or infant brain. The motor impairment may be accompanied by a seizure disorder and by impairment of sensation, cognition, communication and/or behaviour." but also the definition published in 1992 (1) is still in use and probably is the best concise: "Cerebral palsy is an umbrella term covering a group of non-progressive, but often changing, motor impairment syndromes secondary to lesions or anomalies of the brain arising in the early stages of its development".

### 1. 2. Treatment

How to treat person with cerebral palsy? Having in mind a broad spectrum of differences in age of persons at which firm diagnosis is made, inclusions and exclusions criteria, denominator for clinical features (topographical or other classification), and especially what a cause and timing of insult, size and location of lesion, maturational state of system injured, integrity of areas surrounding and contra lateral to lesion, genetic and environmental background, gene-gene and gene-environment interactions, and intervening conditions (timing of start, quality, duration and intensity) was and that we do not know what the potentials of the developing brain of the individual child to adapt to and to compensate for the damage are. It is obvious that there are currently no evidence-based strategies for preventing cerebral palsy and to say which method of treatment is superior. In addition to this fact we should find a balance between physiotherapy (among them nowadays the most used Neurodevelopment treatment– Bobath,

\*Correspondence: Milivoj Velickovic Perat, MD, PhD  
Academy of Developmental Medicine  
Drazgoska 7  
SI-4000 Kranj  
Slovenia  
Phone/fax: +386-4-2361400  
E-mail: milivoj.velickovic@mf.uni-lj.si  
<http://en.perat.eu>

Vojta, Conductive education and Constrain induced movement therapy), speech therapy and occupational therapy and many other approaches used nowadays in the world.

Oral pharmacotherapy (Benzodiazepine, Dantrolene, Baclofen, and many others), Neurosurgical approaches, Local anaesthetics and Local neurolytics (Alcohol, Phenol, Botulinum toxin, Baclofen pump, Selective dorsal rhizotomy, Peripheral neurectomy, ...), Electrical stimulation (Peripheral as is Functional Electrical Stimulation (FES) and Central as are Deep brain stimulation and Dorsal column electrical stimulation), Orthoses (splints, callipers, braces, supports, trusses, casts), Stretching regime to maintain joint range, Surgery for muscle length and bone misalignments, just to mention some of them. Beside this, there are not many proven therapies (Biofeedback, Functional Motor Learning, Acupuncture, the "Adeli" suit, Hyperbaric oxygenation, Homeopathy, Bio-energy, and others).

We should always take in account also Complementary (CAM) and Alternative Treatment (Definition of it is: A group of diverse medical and health care systems, practices, and products that are not presently considered to be part of conventional medicine. The list of what is considered to be CAM changes continually, as those therapies that are proven to be safe and effective become adopted into conventional health care and as new approaches to health care emerge) and a Placebo effect.

A new hope for the brain disorders in the 21st century? Stem cells - to repopulate brain, Transplantation? Is it realistic? Can we use source of donor cells for transplantation, directed at replacement of a specific cell type lost in neurodegenerative diseases, vectors for gene therapy, not only for genetic disorders, but also for replacing specific therapeutic gene products that might help repair brain tissue, and endogenous source for cell replacement. In addition, once transplanted, stem cells have been shown to survive, migrate, and differentiate. Indeed, there is now ample evidence that stem cells exist in the central nervous system throughout life, and the progeny of these stem

cells may have the ability to assume the functional role of neural cells that have been lost.

For the most part, there are key questions that are yet to be answered through well-designed scientific studies

## 2. Conclusion

Advances in medical and surgical management are accelerating at a rapid pace. The only one "best" treatment for all problems of people with cerebral palsy does not exist, but there is no doubt that if we start to treat a problem early on, we will achieve better results. The Early treatment of an infant with abnormal neurological signs is not the same as a treatment of different signs and symptoms of cerebral palsy! Ideally we should start with therapeutic intervention before abnormal movement patterns become dominant and habitual.

Treatment of cerebral palsy must be individualized to each child's needs and must involve parents to teach them effective approaches to apply at home. It should be a holistic approach and we should look on quality of movement patterns and especially on quality of life of a person we are treating.

Progress in the treatment of children and adults with cerebral palsy is a task of exceptional human value and we should do our best to minimize their problems. Brain function is shaped by the interaction of nature, nurture and human brain is especially designed to be influenced by signals from the environment.

We should follow principles of up-to-date therapeutic approaches in the field of 'Evidence based medicine' but we should respect also the traditional and every day approaches used when advanced services provided in developing countries are not available. Dogma is a dictatorship of ideology; especially in treatment approaches. It should be omitted. But wrong ideas are worse than ignorance, too!

## References

1. Mutch LW, Alberman E, Hagberg B, Kodama K, Velickovic MV. Cerebral palsy epidemiology: where are we now and where are we going? *Developmental Medicine and Child Neurology* 1992; 34: 547-555.