

Calcium Hydroxide Treatment Does Not Alter the Susceptibility of *Enterococcus faecalis* Biofilms to Sodium Hypochlorite

Suzette V. van der WAAL, Johannes J. de SOET, Paul R. WESSELINK, Wim CRIELAARD

ABSTRACT

Objective: To investigate the influence of calcium hydroxide ($\text{Ca}(\text{OH})_2$) on susceptibility to disinfection with sodium hypochlorite (NaOCl) of biofilm bacteria.

Methods: Monospecies biofilms of eight *Enterococcus faecalis* strains were subjected to a 2-h challenge with $\text{Ca}(\text{OH})_2$. After a recovery phase, the biofilms were treated with a concentration of NaOCl that was lower than the minimum inhibitory concentration. In a metabolic assay, the efficacy of NaOCl disinfection in $\text{Ca}(\text{OH})_2$ -challenged biofilms and unchallenged biofilms was evaluated. The data were analyzed with Mann-Whitney U and Kruskal-Wallis tests. A P value of less than 0.05 was considered statistically significant.

Results: There were marginal differences in susceptibility to NaOCl among the *E. faecalis* strains. After the $\text{Ca}(\text{OH})_2$ challenge, seven strains remained equally susceptible to NaOCl disinfection whereas one strain became more resistant to NaOCl ($P=0.03$).

Conclusion: After a $\text{Ca}(\text{OH})_2$ challenge, in general *E. faecalis* remained equally susceptible to disinfection with NaOCl .

Keywords: Biofilm, calcium hydroxide, disinfection, endodontic

Please cite this article as "van der Waal S, de Soet JJ, Wesselink PR, Crielaard W. Calcium Hydroxide Treatment Does Not Alter the Susceptibility of *Enterococcus faecalis* Biofilms to Sodium Hypochlorite. Eur Endod J 2017; 2: 30"

From the Department of Preventive Dentistry (S.V.W. ✉ s.vd.waal@acta.nl, J.J.S., W.C.), Academic Centre for Dentistry Amsterdam (ACTA), Amsterdam, The Netherlands; Department of Cariology (S.V.W., P.R.W.), Academic Centre for Dentistry Amsterdam (ACTA), Amsterdam, The Netherlands. Address for Correspondence: Suzette V. van der Waal, Department of Preventive Dentistry, Academic Centre for Dentistry Amsterdam (ACTA), Amsterdam, The Netherlands.

Received 10 April 2017, revision requested 4 May 2017, last revision received 22 July 2017, accepted 31 August 2017.

Published online: 29 November 2017
DOI 10.5152/eej.2017.17022

HIGHLIGHTS

- *Enterococcus faecalis* is not genetically affected by calcium hydroxide.
- Calcium hydroxide does not induce resistance of *Enterococcus faecalis* against disinfection substances such as sodium hypochlorite.
- It is expected that, after the application calcium hydroxide as a root canal dressing, bacteria can still be disinfected with sodium hypochlorite.

INTRODUCTION

Biofilm infection of the root canal system (RCS) accounts for almost all cases of apical periodontitis. Biofilms are complex aggregates of microorganisms attached to a surface and in which the microorganisms are embedded in a matrix of self-produced extracellular polymeric substances. In most biofilms, the matrix consists of over 90% of the dry mass. The matrix protects organisms against desiccation, biocides, some antibiotics, host

immune defenses, etc. (1). These biofilms play an essential role in the creation and maintenance of periapical lesions by both the production of products that are responsible for tissue breakdown and the stimulation of the immune system resulting in an inflammation that needs to be treated.

An important aspect of root canal treatment is to disinfect the RCS by removal of infected tissue using rotating instruments followed by irrigation of a disinfecting agent such as sodium hypochlorite (NaOCl). It is well-known, however, that bacteria remain in the canal space after this instrumentation and irrigation. Thus, many clinicians use a calcium hydroxide ($\text{Ca}(\text{OH})_2$) paste as a temporary root canal dressing to gain additional disinfection and to prevent regrowth of remaining bacteria in between two treatment sessions (2, 3).

The use of $\text{Ca}(\text{OH})_2$ is somewhat questionable. Bacteria can still be recovered from root canals after endodontic treatment with $\text{Ca}(\text{OH})_2$ (4). Also, in *in-vitro* biofilm studies, the disinfecting efficacy

of Ca(OH)₂ seems to be limited (5, 6). Moreover, the complex anatomy of the RCS hinders disinfecting agents from penetrating the entire pulpal space and in a histological study residual biofilms were found in niches of the RCS after Ca(OH)₂ (7). Therefore, it is interesting to study the effect of Ca(OH)₂ on residual biofilms.

Previously, it was shown that Ca(OH)₂ causes changes in the configuration of the biofilm matrix (6, 8). *Enterococcus faecalis* has been associated with failed endodontic treatments (9). One study reported that *E. faecalis* still formed biofilms in the presence of Ca(OH)₂ and that the bacteria were encompassed in an even denser matrix. Moreover, approximately 90% of these cells were viable (8). Another study showed that a dense flocculation occurred in dual-species biofilms of *E. faecalis* and *Pseudomonas aeruginosa* after a treatment with Ca(OH)₂. Also, there was only a factor 100 reduction of viable cells in these biofilms, which contained on average 3×10^8 cells (6). It has also been reported that *E. faecalis* can produce more biofilm mass and protein in response to high pH or sub-inhibitory doses of antimicrobials (10, 11).

From the literature we know that calcium, which is released when Ca(OH)₂ is solubilized in water, stimulates the formation of extracellular polysaccharides (EPS) and that it increases the yield strength of biofilms (12, 13). Plus, a high pH favors the uptake of calcium in the EPS (14).

The above-mentioned *in-vitro* observations combined with the clinical findings support the theory that Ca(OH)₂ may stimulate the formation of a more dense biofilm of *E. faecalis* that may be less sensitive to disinfection in a subsequent treatment session. The formation of a dense matrix or an increase in the coherence of the biofilm matrix can protect the bacteria from disinfecting measures. Also, the genetic make-up of the bacteria may be altered by Ca(OH)₂ (15). Until now, no literature is available on inadequate Ca(OH)₂ treatments and the possible effects on the susceptibility to disinfection of the remaining biofilm bacteria. Therefore, the aim of this study was to investigate the influence of calcium hydroxide (Ca(OH)₂) on the susceptibility of biofilm bacteria to disinfection with sodium hypochlorite. The null-hypothesis of this study was that Ca(OH)₂ treatment of *E. faecalis* strains in an active attachment biofilm model does not induce changes in the sensitivity of these biofilms to NaOCl resulting in a more resistant biofilm.

METHODS

Approval

The ethical committee of Academic Centre Dentistry Amsterdam, ACTA, confirmed that the Medical Research Involving Human Subjects Act (WMO) did not apply to this study and that this study was performed according to the ethical guidelines at ACTA. The reference number is 2017036Waal.

TABLE 1. Origins and sources of the employed *E. faecalis* strains.

Strain name	Origin	Source	Reference
E1	oral rinse	Sedgley	(18, 27)
E2	oral rinse	Sedgley	(18, 27)
E3	oral rinse	Sedgley	(18, 27)
AA-OR34	oral rinse	Sedgley	(18, 28)
ER3/2s	orthograde retreatment	Sedgley	(18)
ER5/1	orthograde retreatment	Sedgley	(18)
OS16	oral rinse	Sedgley	(18, 29)
V583	Lab strain	Sahm	(30)

Medium

Biofilms were grown in modified semi-defined broth, pH 7.1, with 0.2% sucrose (BMS) whose composition has been previously published (16).

Biofilm Growth, Treatment and Resazurin metabolism

The resazurin metabolism assay and the model have been described previously (17). It consists of a standard 96-well microtiter plate and a lid with 96 polystyrene pegs that fit into the wells (TSP; Nunc, Roskilde, Denmark).

The origins of the *E. faecalis* strains can be found in Table 1. *E. faecalis* strains E1, E2, E3, AA-OR34 and OS16 originated from oral rinses and were kindly provided by Christine Sedgley (18). Strains ER3/2s and ER5/1, which have not been published before, also originated from the Sedgley lab and were isolated from orthograde retreatment. Strain V583 (ATCC700802) was used as reference strain. All strains were maintained in stocks at -80°C, and fresh cultures were obtained by the addition of 500 µL of frozen stock culture to 5 mL semi-defined medium supplemented with yeast +0.36% glucose and incubation of 6 h in an anaerobic jar (80% N₂, 10% H₂ and 10% CO₂) at 37°C (16). By initial optical densities at 600 nm (OD₆₀₀) at culture density 0.8 (Spectramax M2, Sunnyvale, CA, USA) the cultures were in a late log phase and contained approximately 5×10^8 cells mL⁻¹. After 1:100 dilution with BMS, 200 µL of this inoculant was dispersed per well of a 96-well microtiter plate and the plate was covered with the TSP lid. The medium was refreshed after 18 h. After 24 h, the TSP lid with adhered biofilms was transferred to a 96-well plate with 200 µL well⁻¹ BMS or 10% Ca(OH)₂ (E. Merck, Darmstadt, Germany) (weight/volume, w/v) in BMS (slurry), pH 12.6 for 2 h. Subsequently, the biofilms were washed twice in 200 µL phosphate buffered saline (PBS) and transferred to BMS for an overnight recovery of the biofilms at 37°C anaerobic. Shortly before the final treatments, the free available chlorine concentration of the NaOCl (Orphi Farma, Lage Zwaluwe, The Netherlands) was verified with an iodometric titration procedure and NaOCl was diluted in demineralized water to obtain 1 or 10 ppm solutions of active chlorine (19). 1 ppm NaOCl appeared to be a sub-bactericidal dose for these particular biofilms and therefore suitable for detecting a possible difference between the groups. When the bio-




Figure 1. The treatment scheme of the viability assay in which biofilms were exposed to calcium hydroxide ($\text{Ca}(\text{OH})_2$) and were subsequently treated with sodium hypochlorite (NaOCl)




Figure 2. Metabolic activity of various *Enterococcus faecalis* strains with or without pre-treatment with calcium hydroxide and after a 5-min treatment with 1 or 10 ppm sodium hypochlorite. The controls were treated with phosphate buffered saline. Only one strain, *E. faecalis* OS16 (marked with a star), became less susceptible to NaOCl after a pretreatment with $\text{Ca}(\text{OH})_2$ ($P=0.03$). Error bars represent standard error of the mean. The individual strains are arranged from least susceptible to NaOCl (left) to most susceptible to NaOCl (right). Differences in susceptibility between strains is shown with the bars depicted above the graph.

films were 42-h old, they were treated for 5 min in a 96 wells plate that contained 1 or 10 ppm NaOCl 210 μL in each well. Then, NaOCl was neutralized for 5 min with 1% sodium thio-sulphate (Merck) in 2% buffered peptone water (BPW; Oxoid, Basingstoke, UK) after which the biofilms were washed twice in PBS. The negative control group was treated with PBS. Sterile pegs were included to determine the background measurement. Subsequently, the pegs were immersed in 0.016 mg mL^{-1} resazurin (Sigma-Aldrich, St. Louis, MO, USA) in BMS and incubated at 37°C in air. Metabolic activity was determined by measuring the fluorescence intensity (FI) of each well, which was recorded at room temperature in a flu-

orimeter (Spectramax M2; Molecular Devices, Sunnyvale, CA) using 485-nm excitation and 580-nm emission wavelengths. Readings were taken at 3 h. For an overview of the culturing and handling of the biofilms see Figure 1. Each experiment was performed in duplo and the assay was repeated on two separate occasions. The average background measurement was subtracted from the FI readings, which were then averaged.

Statistical analysis

All data were analysed using statistical analysis software (SPSS Version 21, IBM Corp.; Armonk, NY, USA). The data were not normally distributed and therefore were non-parametrically tested. Mann Whitney U tests were used to calculate differences between groups, while a Kruskal-Wallis test was used to test the median of multiple groups. A P value of less than 0.05 was considered significant.

RESULTS

There were significant differences in the susceptibility to NaOCl between the individual strains of *E. faecalis*: strain ERS/1 was the least susceptible, whereas AA-OR34 was the most susceptible (Figure 2). These differences were independent of the use of $\text{Ca}(\text{OH})_2$.

On the whole, *E. faecalis* biofilms that had been exposed to $\text{Ca}(\text{OH})_2$ were equally susceptible to NaOCl disinfection as the control biofilms ($P=0.343$). However, when evaluating the strains individually, *E. faecalis* OS16 was less susceptible to NaOCl disinfection ($P=0.03$) (Figure 2).

DISCUSSION

The aim of the present study was to investigate whether susceptibility to disinfection by NaOCl of *E. faecalis* biofilms would change after treatment with $\text{Ca}(\text{OH})_2$. The current results show that, after $\text{Ca}(\text{OH})_2$ treatment, the majority of the tested strains remained equally susceptible to disinfection with NaOCl. One strain, OS16, became significantly less susceptible and none of the strains became more susceptible to NaOCl disinfection. The reference strain, V583, was not different from the oral clinical isolates. The null-hypothesis was therefore accepted.

Although, the current model does not resemble the clinical situation, it was specifically designed to investigate our fundamental question about the behavior of *E. faecalis* after $\text{Ca}(\text{OH})_2$ treatment. *E. faecalis* is a bacterial species often encountered as a recalcitrant organism in unsuccessful root canal treatments. It is well-studied in laboratory experiments and, because of its tenacity, is often used as a target organism in disinfection experiments. It is known to survive high pH (6, 8, 10, 20). The presence of calcium stimulates the formation of EPS in *E. faecalis* biofilms and this may help bacteria within these biofilms to survive certain antimicrobial conditions (12).

The biofilms were grown on an inert substrate (polystyrene pegs) to exclude any possible interference of dentine. Low sub-inhibitory concentrations of NaOCl were used because the susceptibility of *E. faecalis* before and after Ca(OH)₂ application cannot be measured with clinically applicable NaOCl concentrations. Prior to the current study, sub-bactericidal concentrations and incubation-time of Ca(OH)₂ and NaOCl were established for these *E. faecalis* biofilms. In a previous study, a saturated slurry of 10% Ca(OH)₂ at pH 12.5 was shown to be a non-inhibitory concentration in biofilms. Since the solubility constant of Ca(OH)₂ at room temperature is 1.85 g L⁻¹, an active concentration of approx. 0.2% OH⁻ can be obtained. Commercial Ca(OH)₂ pastes/slurries are sold at higher concentrations (30-35%) but will result in a similarly saturated active concentration of 0.2%. The sub-inhibitory treatment of NaOCl was determined at 5 min 1 ppm, which appeared to be a little higher than previously found in similar biofilms (21). If this model is to be employed for investigating the susceptibility of dual- or multi-species biofilms, the concentration of NaOCl will need to be adjusted depending on the type of species and the age of the biofilms (22). It can be argued that these low concentrations of NaOCl are not representative of the clinical situation, but since compounds such as NaOCl strongly react with tissues within the endodontic canals, these low concentrations are likely to be found in the more remote sites of the RCS. This may be one of the reasons that bacteria survive endodontic treatment.

By using a high-throughput system with resazurin, the number of viable cells for the NaOCl sensitivity assay should be roughly similar. This was one of the limitations of the many studies on antimicrobial properties of Ca(OH)₂: variation in inoculum size (23). It was therefore necessary to recover the biofilms for 16 hours after the Ca(OH)₂ treatment so that the bacterial numbers could be restored to the same number of bacteria in the control biofilms (data not published). A previous study demonstrated that, with Ca(OH)₂, the biofilms stayed attached to the surface (6). Therefore, detachment after Ca(OH)₂ application, resulting in a lower resazurin signal, was not observed. Also, there was no difference between the negative controls of the non-Ca(OH)₂ and Ca(OH)₂ groups. Consequently, there were no indications that the results from this study were biased by experimental errors.

Calcium hydroxide is a chemical compound that has one calcium ion and two hydroxide ions (OH⁻). These OH⁻ ions render the environment alkaline in which many bacteria cannot survive. The presence of the free OH⁻ ion as a radical is assumed to have direct effects on the bacterial membrane, protein denaturation and bacterial DNA damage. Certain issues of Ca(OH)₂, however, have been largely overlooked. OH⁻ ions are reducing molecules and are unstable in a solution as they tend to react with CO₂ or other oxides (24). When OH⁻ is consumed, Ca(OH)₂ solubilises and, besides OH⁻, Ca²⁺ ions are released. Ca²⁺ ions are divalent cations and as such they bind to the negatively charged surfaces of bacteria, to dentin and

to the EPS of the biofilm matrix. This series of events can explain why (traces of) Ca(OH)₂ are difficult to remove from the RCS after use. The solubilisation of the depot of Ca(OH)₂ in the slurry or paste is a continuous process until the depot has been depleted.

This study was designed for *E. faecalis*. Besides being associated with failed endodontic treatment, *E. faecalis* is easy to culture *in vitro* and therefore is a suitable test microorganism (9). Many properties of *E. faecalis* have already been investigated. For example, a study of the antimicrobial susceptibility of enterococci isolated from root canal showed that *E. faecalis* can be highly resistant to certain antibiotics (25). Our study, however, is the first to show that, despite the changes that occur in the biofilms after the application of Ca(OH)₂, *E. faecalis* in general does not become more resistant to disinfection by NaOCl. We should realise though, that the present study was performed on mono-species biofilms and it may be that multispecies interaction, which occurs in endodontic biofilms, can result in more resistance to either Ca(OH)₂ or NaOCl (26). It would be interesting to see, in future studies, if our theory holds true when endodontic microcosm biofilms are used. Such a study would possibly also give information about the changes in bacterial composition of the biofilms after the application of Ca(OH)₂.

The current findings are clinically relevant because, after the application of Ca(OH)₂, there are often residual bacteria. Also, *E. faecalis* is often found in such secondary infections. This study has, however, demonstrated that the implications of an inefficient treatment with Ca(OH)₂ may be minor.

CONCLUSION

The current results indicate that, after a challenge with Ca(OH)₂, one of eight *E. faecalis* strains became less susceptible to disinfection. However, when using a higher dose of 10 ppm NaOCl, this strain was also inactivated. Seven of eight strains of *E. faecalis* remained equally susceptible to disinfection with NaOCl. Therefore, it appears that, in general, Ca(OH)₂ does not affect the susceptibility of *E. faecalis* to NaOCl.

Ethics Committee Approval: Ethics committee approval was received for this study from the ethics committee of the Academic Centre Dentistry Amsterdam (ACTA) / 2017036Waal.

Informed Consent: Not applicable as this was a microbiology study with laboratory strains.

Peer-review: Externally peer-reviewed.

Author Contributions: Concept - S.W., P.W., W.C., J.S.; Design - S.W.; Supervision - W.C., J.S.; Resources - W.C.; Materials - W.C.; Data Collection and/or Processing - S.W.; Analysis and/or Interpretation - S.W., J.S.; Literature Search - S.W., P.W.; Writing Manuscript - S.W.; Critical Review - P.W., W.C., J.S.

Acknowledgements: The authors wish to thank Christine Sedgley for providing the clinical isolates of *E. faecalis*.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study has been supported by ACTA.

REFERENCES

1. Flemming HC, Wingender J. The biofilm matrix. *Nat Rev Microbiol* 2010; 8(9): 623-33. [\[CrossRef\]](#)
2. Palmer NO, Ahmed M, Grieveson B. An investigation of current endodontic practice and training needs in primary care in the north west of England. *Br Dent J* 2009; 206(11): 584-5. [\[CrossRef\]](#)
3. Savani GM, Sabbah W, Sedgley CM, Whitten B. Current trends in endodontic treatment by general dental practitioners: report of a United States national survey. *J Endod* 2014; 40(5): 618-24. [\[CrossRef\]](#)
4. Sathorn C, Parashos P, Messer HH. How useful is root canal culturing in predicting treatment outcome? *J Endod* 2007; 33(3): 220-5. [\[CrossRef\]](#)
5. Abdullah M, Ng YL, Gulabivala K, Moles DR, Spratt DA. Susceptibilities of two *Enterococcus faecalis* phenotypes to root canal medications. *J Endod* 2005; 31(1): 30-6. [\[CrossRef\]](#)
6. van der Waal SV, van der Sluis LW, Özok AR, Exterkate RA, van Marle J, Wesselink PR, et al. The effects of hyperosmosis or high pH on a dual-species biofilm of *Enterococcus faecalis* and *Pseudomonas aeruginosa*: an in vitro study. *Int Endod J* 2011; 44(12): 1110-7. [\[CrossRef\]](#)
7. Vera J, Siqueira Jr. JF, Ricucci D, Loghin S, Fernández N, Flores B, et al. One-versus two-visit endodontic treatment of teeth with apical periodontitis: a histobacteriologic study. *J Endod* 2012; 38(8): 1040-52. [\[CrossRef\]](#)
8. Distel JW, Hatton JF, Gillespie MJ. Biofilm formation in medicated root canals. *J Endod* 2002; 28(10): 689-93. [\[CrossRef\]](#)
9. Chávez De Paz LE, Dahlén G, Molander A, Möller A, Bergenholtz G. Bacteria recovered from teeth with apical periodontitis after antimicrobial endodontic treatment. *Int Endod J* 2003; 36(7): 500-8. [\[CrossRef\]](#)
10. Chávez de Paz LE, Bergenholtz G, Dahlén G, Svensäter G. Response to alkaline stress by root canal bacteria in biofilms. *Int Endod J* 2007; 40(5): 344-55. [\[CrossRef\]](#)
11. Wilson CE, Cathro PC, Rogers AH, Briggs N, Zilm PS. Clonal diversity in biofilm formation by *Enterococcus faecalis* in response to environmental stress associated with endodontic irrigants and medicaments. *Int Endod J* 2015; 48(3): 210-19. [\[CrossRef\]](#)
12. Safari A, Habimana O, Allen A, Casey E. The significance of calcium ions on *Pseudomonas fluorescens* biofilms - a structural and mechanical study. *Biofouling* 2014; 30(7): 859-69. [\[CrossRef\]](#)
13. Körstgens V, Flemming HC, Wingender J, Borchard W. Influence of calcium ions on the mechanical properties of a model biofilm of mucoid *Pseudomonas aeruginosa*. *Water Sci Technol : J Int Ass Water Pollution Res* 2001; 43(6): 49-57.
14. Geddie JL, Sutherland IW. Uptake of Metals by Bacterial Polysaccharides. *J Appl Bacteriol* 1993; 74(4): 467-72. [\[CrossRef\]](#)
15. Kohanski MA, DePristo MA, Collins JJ. Sublethal antibiotic treatment leads to multidrug resistance via radical-induced mutagenesis. *Molecular Cell* 2010; 37(3): 311-20. [\[CrossRef\]](#)
16. Deng DM, Hoogenkamp MA, Exterkate RA, Jiang LM, van der Sluis LW, Ten Cate JM, et al. Influence of *Streptococcus mutans* on *Enterococcus faecalis* biofilm formation. *J Endod* 2009; 35(9): 1249-52. [\[CrossRef\]](#)
17. Jiang LM, Hoogenkamp MA, van der Sluis LW, Wesselink PR, Crielaard W, Deng DM. Resazurin metabolism assay for root canal disinfectant evaluation on dual-species biofilms. *J Endod* 2011; 37(1): 31-5. [\[CrossRef\]](#)
18. Duggan JM, Sedgley CM. Biofilm formation of oral and endodontic *Enterococcus faecalis*. *J Endod* 2007; 33(7): 815-8. [\[CrossRef\]](#)
19. Vogel A. A textbook of quantitative inorganic analysis. 3rd ed. London, UK: Longmans; 1962.
20. Nakajo K, Komori R, Ishikawa S, Ueno T, Suzuki Y, Iwami Y, et al. Resistance to acidic and alkaline environments in the endodontic pathogen *Enterococcus faecalis*. *Oral Microbiol Immunol* 2006; 21(5): 283-8. [\[CrossRef\]](#)
21. Arias-Moliz MT, Ferrer-Luque CM, Espigares-García M, Baca P. *Enterococcus faecalis* biofilms eradication by root canal irrigants. *J Endod* 2009; 35(5): 711-4. [\[CrossRef\]](#)
22. Özok AR, Wu MK, Luppens SB, Wesselink PR. Comparison of growth and susceptibility to sodium hypochlorite of mono- and dual-species biofilms of *Fusobacterium nucleatum* and *Peptostreptococcus micromonas*. *J Endod* 2007; 33(7): 819-22. [\[CrossRef\]](#)
23. Kim D, Kim E. Antimicrobial effect of calcium hydroxide as an intracanal medicament in root canal treatment: a literature review - Part I. In vitro studies. *Rest Dent & Endod* 2014; 39(4): 241-52. [\[CrossRef\]](#)
24. Yoo M, Han SJ, Wee JH. Carbon dioxide capture capacity of sodium hydroxide aqueous solution. *J Environ Manage* 2013; 114: 512-9. [\[CrossRef\]](#)
25. Dahlén G, Samuelsson W, Molander A, Reit C. Identification and antimicrobial susceptibility of enterococci isolated from the root canal. *Oral Microbiol Immunol* 2000; 15(5): 309-12. [\[CrossRef\]](#)
26. Özok AR, Persoon IF, Huse SM, Keijser BJB, Wesselink PR, Crielaard W, et al. Ecology of the microbiome of the infected root canal system: a comparison between apical and coronal root segments. *Int Endod J* 2012; 45(6): 530-41. [\[CrossRef\]](#)
27. Sedgley CM, Lennan SL, Clewell DB. Prevalence, phenotype and genotype of oral enterococci. *Oral Microbiol Immunol* 2004; 19(2): 95-101. [\[CrossRef\]](#)
28. Sedgley CM, Nagel AC, Shelburne CE, Clewell DB, Appelbe O, Molander A. Quantitative real-time PCR detection of oral *Enterococcus faecalis* in humans. *Arch Oral Biol* 2005; 50(6): 575-83. [\[CrossRef\]](#)
29. Sedgley C, Buck G, Appelbe O. Prevalence of *Enterococcus faecalis* at multiple oral sites in endodontic patients using culture and PCR. *J Endod* 2006; 32(2): 104-9. [\[CrossRef\]](#)
30. Sahm DF, Kissinger J, Gilmore MS, Murray PR, Mulder R, Solliday J, et al. In vitro susceptibility studies of vancomycin-resistant *Enterococcus faecalis*. *Antimicrobiol Agents Chemother* 1989; 33(9): 1588-91. [\[CrossRef\]](#)