

Effect of exercise doses on functional recovery in neonatal brachial plexus palsy: A randomized controlled study

 Nilay Sahin,¹  Ali Yavuz Karahan²

¹Department of Physical Therapy and Rehabilitation, Balikesir University Faculty of Medicine, Balikesir, Turkey

²Department of Physical Therapy and Rehabilitation, Usak University Faculty of Medicine, Usak, Turkey

ABSTRACT

OBJECTIVE: The aim of this study was to examine how much recovery upper obstetric brachial plexus palsy (OBPP) patients showed with exercises when they were 3, 6 and 12 months old; and to evaluate whether the exercise treatment given at different frequencies contribute to this recovery or not.

METHODS: This study was designed as a randomized controlled trial. Sixty cases that were referred to Pediatric Rehabilitation and Pediatric Orthopedic Clinics with the diagnosis of having Group I and II OBPP according to Narakas classification were included in the study. Subjects were randomly divided into two treatment groups. The first group had intense exercise program three times daily, and the second group had a standard exercise program once in a day. The subjects were assessed using a passive-active range of motion (ROM) and Hospital for Sick Children muscle grading system at their first clinic visit and every month after until they became 12 months old.

RESULTS: In both groups, a significant recovery was observed in ROM and muscle strength of all movements of the shoulder, elbow flexion, and forearm supination at the 3rd, 6th, and 12nd-month reassessments, whereas a significant difference was not achieved on both parameters between two groups.

CONCLUSION: According to the results exercise frequency did not affect recovery rate and results in the cases with OBPP and exercises were influential against possible complications that may occur.

Keywords: Brachial plexus lesion; exercise; functional recovery; muscle strength.

Cite this article as: Sahin N., Karahan A. Y. Effect of exercise doses on functional recovery in neonatal brachial plexus palsy: A randomized controlled study.

Obstetric brachial plexus palsy (OBPP) is a flaccid paralysis mostly seen on arm due to the injury of brachial plexus during birth. It occurs as a result of distention and tear in the body of brachial plexus and avulsion of spinal roots [1, 2]. At every 1000 birth, OBPP incidence is 0.3–1.56%, recovery rate is 84% and permanent damage rate is 0.5–25%. So OBPP is an important neonatal morbidity cause [3–5]. The most observed form of OBPP is the involvement of upper brachial plexus (C5-6,7). In C5-6 injuries, shoulder abductors,

internal rotators, and forearm flexor muscles are mostly affected, the decline in forearm supination movement can be seen, but hand functions are retained. If an injury to C7 is added than triceps and wrist extensors are also affected [4, 6, 7].

Although an enough spontaneous recovery to use extremity functionally can be gained in most of the upper OBPP patients, all these patients are given an exercise program at the first stage [2, 8–10]. The aim to give exercises is mostly to prevent contracture formation and to

Received: July 08, 2017 *Accepted:* December 28, 2017 *Online:* August 07, 2018



Correspondence: Dr. Ali Yavuz KARAHAN. Usak Universitesi Tip Fakultesi, Fiziksel Tedavi ve Rehabilitasyon Anabilim Dalı, 64200 Usak, Turkey.
Tel: +90 276 221 22 33 e-mail: ayk222@hotmail.com

© Copyright 2018 by Istanbul Provincial Directorate of Health - Available online at www.northclinist.com

avoid muscle imbalance [4, 7]. There is no accurate data on the effect of any particular exercise is more than the other, or one has additional effects on recovery than others to prevent complication; besides there is no consensus on what frequency, intensity or duration should exercise therapy be given. While some authors claimed that exercises do not have any effect on the recovery of the patients, others suggested that exercises could contribute to recovery by positively affecting regenerative process [4].

It was hypothesized that intense exercise programs given in cases with peripheral nerve or spinal cord injury may accelerate recovery by contributing to plasticity development [11–13]. When no improvement is gained in biceps muscle function in patients with OBPP in the first three months during follow-up, shoulder dysfunction or residual upper extremity incapability often develops. Therefore, some researchers recommend going further to surgical interventions in cases that recovery is not observed in biceps muscle function within three to six months [9]. Therefore, the aim of this study was to examine how much recovery upper obstetric brachial plexus palsy (OBPP) patients showed with exercises when they were 3, 6 and 12 months old; and to evaluate whether the exercise treatment given at different frequencies contribute to this recovery or not.

MATERIALS AND METHODS

This study conducted in physical medicine and rehabilitation department of Balıkesir University between June and December 2014. The study was approved by the local research ethics committee of Selçuk University's Faculty of Medicine (No: 276/2014). Written informed consent was obtained from parents of all patients for being included in the study. Sixty cases that were referred to Pediatric Rehabilitation and Pediatric Orthopedic Clinics with the diagnosis of group I and II OBPP according to Narakas classification, were included in the study following a multidisciplinary approach; physical examination and electromyography (EMG) assessments. The patients in group III and IV according to Narakas classification 2, the ones who had another comorbid neurological problem, contracture and/or fracture on the affected arm were excluded from the study. The parents of patients were elucidated with precise information about the study and all of them signed approval forms.

Sixty patients were divided into two groups randomly (For the randomization-allocation concealment, all the randomization numbers were concealed in separate en-

velopes and marked by the patient number on the outer envelope. For randomization, the patients admitted into the program were divided into groups); the first group had intense exercise program three times daily (intensive exercise group [IEG] as a group I), and the second group defined as a control group and had a standard exercise program once in a day (exercise group [EG] as group II).

Exercise program: Passive ROM exercises were given to all joints of the upper extremities at first, and in the following months active ROM and muscle enhancement exercises (according to the cooperation level of the children) were started. The duration time of exercises were related to the children's cooperation. Because biceps muscle force of 3 patients in the group 1 and 1 patient in the group II was weak, they were excluded from the study. The subjects were assessed using passive-active range of motion (ROM) degrees and Hospital for Sick Children muscle grading system on their first clinic visit and every month after, until they became 12 months old [5].

Assessment Criteria

1. Narakas classification: This classification is composed of four groups: Group I C5-6 shoulder and biceps paralysis, Group II C5-7 paralysis of shoulder, biceps and forearm extensors, Group III C5-Th1 complete paralysis of the extremity and Group IV C5-Th1 complete paralysis of extremity accompanying Horner syndrome [5].

2. Range of motion: ROM was evaluated both passively and actively with joint specific movements: shoulder ROM: flexion, abduction, extension, external/internal rotations; elbow ROM: flexion, extension, supination and pronation; wrist ROM: extension and flexion. Active ROM was measured on three different days following each other and calculated by averaging these measures by the same observer, which proved intra-observer reliability.

3. Muscle testing: Shoulder flexors and abductors, elbow flexor and forearm supinator muscles were evaluated using Hospital for Sick Children muscle grading system (Clarke and Curtis). This classification system is appropriate for children younger than 3 years old, which measures all joints without any commands. The Hospital for Sick Children scale is not a 'muscle grading system' but is an assessment of spontaneous active movement. There are 7 stages totally, in which the gravity is eliminated until stage 4 (Table 1).

Physiotherapy program: The families were educated in the hospital by a talented physiotherapist with a five-

TABLE 1. Hospital for sick children muscle grading system

Observation	Muscle grade
Gravity eliminated	0
No contraction	1
Motion < ½ range	2
Motion >½ range	3
Full motion	4
Against gravity	
Motion < ½ range	5
Motion >½ range	6
Full motion	7

day oral, visual and applied training program, in which the parents were than tested. The patients were examined and exercise programs were checked by the same physician every month until they became 12 months old. It was ordered to make the exercises three times daily for the IEG and once daily for the EG.

Statistical Analysis

Statistical evaluation was performed using SPSS 16 (SPSS Inc. Released 2007. SPSS for Windows, Version 16.0. Chicago, SPSS Inc.). The one-sample Kolmogorov-Smirnov test was applied for non-parametric values that comply with a normal distribution in inner-group assessments. Analysis of variance (ANOVA) was used in two-way repeated (time and groups) measurements to compare data from the parameters repeatedly measured in the inner group. In cases in which the variance analysis test result was significant, Bonferroni correction was performed to compare between groups as a post hoc test. The Bonferroni Student t-test was used to detect different groups. The Student t-test was used for independent groups to analyze data that complied with a normal distribution. P values less than 0.05 were considered statistically significant ($p < 0.05$).

RESULTS

Since mean biceps muscle force of 4 patients was 1.42 when they were 4 months old, these children were referred to surgery and excluded from the study. So totally 62 patients (38 boys–24 girls), with a mean age of 7.50 ± 5.18 days were included in the study. The affected

TABLE 2. Characteristics of patients of obstetric brachial plexus palsy

Age (days)	7.50±5.18
Sex (male/female)	38/15
Weight (gr)	4388
Affected side (left/right)	42:20

side was left in 42 cases, and right in 20 cases. All infants were born vaginally and 52 cases had difficult labor history. The average birth weight was 4388 grams (Table 2). 97% of the subjects were in group I while 3% were in group II according to Narakas classification.

Statistical analysis inside the groups: In terms of active ROM, while a statistically significant improvement was gained in both groups between 1st–3rd months, 3rd–6th months and 6th–12nd months in shoulder abduction, external rotation and flexion, elbow flexion and forearm supination and between 1st–3rd months and 3rd–6th months in shoulder internal rotation and extension ($p < 0.05$); there was no significant difference in elbow extension, forearm pronation and wrist extension since they already had nearly normal values at the beginning of the treatment ($p > 0.05$). Active ROM measures of the patients are shown in Table 3. There was a statistically significant improvement in the muscle strength in both groups between 1st–6th months and 6th–12nd months in shoulder abduction, between 1st–3rd and 6th–12nd months in shoulder flexion, and between 1st, 3rd, 6th, and 12nd months in elbow flexion and forearm supination ($p < 0.05$) (Table 4).

Statistical analysis between the groups: No statistically significant improvement was observed between the groups in EMP measures and muscle force assessments. ($p > 0.05$).

DISCUSSION

In upper brachial plexus OBPP patients there was no significant difference between the groups who had intensive exercise program and those who did not. Only 4 of the cases were further referred to surgery. In all the cases except for these 4 cases, good arm functions were observed while they were three months old, and nearly normal functions were found while they were 12 months old. So almost exact recovery was gained in 94% of the cases with OBPP in our study. The patients with OBPP

TABLE 3. Measurements of active range of motion in patients with obstetric brachial plexus palsy

	Birth		3 months		6 months		12 months	
	Group I	Group II	Group I	Group II	Group I	Group II	Group I	Group II
Shoulder (°)								
Abduction	11.92 (4.38)	12.82 (3.63)	38.71 (3.38)*	38.04 (3.61)†	62.92 (2.46)*	62.82 (2.53)†	87.63 (2.53)*	87.40 (2.55)†
External rotation	0	0	18.07 (2.46)*	18.03 (2.24)†	23.92 (2.48)*	23.47 (2.35)†	28.20 (2.42)*	28.04 (2.49)†
Internal rotation	39.74 (3.79)	39.56 (3.60)	60.38 (4.35)*	60.21 (3.19)†	78.33 (2.38)*	77.60 (2.56)†	79.23 (1.82)*	78.96 (2.24)†
Extension	0	0	18.84 (2.13)*	17.60 (2.55)†	22.15 (2.13)*	22.17 (2.53)†	22.53 (2.33)*	22.04 (2.38)†
Flexion	19.87 (3.34)	19.13 (3.88)	51.53 (2.33)*	51.39 (2.55)†	66.53 (2.33)*	65.43 (2.21)†	83.58 (2.27)*	82.60 (2.45)†
Elbow (°)								
Flexion	0	0	68.07 (2.46)*	67.60 (2.55)†	77.92 (2.46)*	77.39 (2.54)†	102.82 (4.55)*	102.78 (5.10)†
Extension	103.07 (4.67)	103.47 (4.86)	115.89 (4.98)*	115.65 (5.06)†	118.46 (3.65)*	117.82 (4.21)†	118.46 (3.65)*	117.82 (4.21)†
Forearm (°)								
Pronation	118.46 (2.33)	118.69 (2.24)	121.38 (5.05)*	121.21 (5.10)†	121.38 (5.05)*	121.21 (5.10)†	121.38 (5.05)*	121.21 (5.10)†
Supination	0	0	36.79 (2.42)*	37.60 (2.55)†	72.79 (2.42)*	72.60 (2.55)†	106.66 (4.77)*	105.21 (5.10)†
Wrist (°)								
	81.79 (2.42)	81.52 (2.35)	88.20 (2.42)*	88.47 (2.35)†	88.20 (2.42)*	88.47 (2.35)†	88.20 (2.42)*	88.47 (2.35)†
Flexion	71.46 (2.33)	71.95 (2.49)	71.46 (2.33)*	71.95 (2.49)†	71.46 (2.33)*	71.95 (2.49)†	71.46 (2.33)*	71.95 (2.49)†

*: Statistically significant differences according to the Birth value inner Group I evaluations ($p < 0.05$). †: Statistically significant differences according to the Birth value inner Group II evaluations ($p < 0.05$). $p > 0.05$ for each between group evaluations; (°): Degree of movement at a joint.

TABLE 4. Hospital for Sick Children muscle grading system in patients with obstetric brachial plexus palsy

	Birth		3 months		6 months		12 months	
	Group I	Group II	Group I	Group II	Group I	Group II	Group I	Group II
Shoulder								
Abduction	3.50 (0.46)	3.52 (0.51)	4.07 (0.57)*	4.01 (0.73)†	5.51 (0.50)*	5.56 (0.50)†	6.64 (0.48)*	6.56 (0.50)†
Flexion	3.61 (0.49)	3.65 (0.48)	5.40 (0.46)*	5.47 (0.51)†	5.43 (0.48)*	5.48 (0.52)†	6.46 (0.42)*	6.30 (0.47)†
Elbow								
Flexion	0.56 (0.50)	0.52 (0.51)	4.71 (0.64)*	4.82 (0.77)†	5.45 (0.64)*	5.43 (0.50)†	6.38 (0.45)*	6.39 (0.49)†
Forearm								
Supination	0.41 (0.50)	0.43 (0.50)	2.58 (0.52)*	2.60(0.49)†	5.53 (0.49)*	5.52 (0.51)†	6.30 (0.40)*	6.34 (0.48)†

*: Statistically significant differences according to the Birth value inner Group I evaluations ($p < 0.05$). †: Statistically significant differences according to the Birth value inner Group II evaluations ($p < 0.05$). $p > 0.05$ for each between group evaluations.

at the C5-6 level, generally display improvement in a period from 3 months to 9 months of age [3–5, 7]. It is reported that improvement period may continue until 12 months in some studies [5].

The diagnosis of OBPP is based on a precise physical examination with a support of EMG study. The valid pa-

rameters used to evaluate improvement in OBPP cases are ROM measurements, EMG studies, and Clarke and Curtis muscle classification system for children under the age of [3–5]. If a common development cannot be gained in the shoulder and biceps muscles while the child is in a period of 3 months and 6 months old, then he/

she is referred to surgery [4, 7]. While 70–95% of the patients improve almost completely, 10% may require surgery [4, 7, 14]. Likely 6% of our subjects had operations, 94% displayed a nearly complete improvement between 6–12 months. The increase in biceps and shoulder muscles showed a rapid incline in the 3rd month and became almost normal in 6th–12nd months.

Physiotherapy program is the recommended treatment option for patients with C5-7 OBPP at the first stage. This conservative treatment includes exercise, splint, and electrical stimulation. The purpose of conservative treatment is mainly to prevent contracture formation, hypoplastic arm development, dislocation of the radial head and negative psychosocial consequences [1, 2, 4, 7]. Although a spontaneous recovery can be seen in most of the cases, contracture formation in muscles, dislocation in shoulder and elbow may prevent functional recovery [7]. At this point, exercise program substantially contributes to the recovery of most of the patients.

However, there is no accurate data about the frequency, intensity and time-planning of exercise programs. It is suggested that permanent disability may develop in some of the cases despite physiotherapy treatment [7]. This situation has raised more questions about the place of exercise in OBPP. Some authors suggested that exercises did not have much effect on recovery [4]. But the other some have defended that intensive exercise program has a positive impact on recovery. However, they have still not clearly explained how this effect occurs [10].

When we examine the processes in the recovery of OBPP; we will observe that not only peripheral reconstruction but also central and spinal cord plasticity play a significant role in OBPP recovery [6, 8, 15]. It is hypothesized that physical exercises increase plasticity especially in the spinal cord and central area by inclining ependymal cell proliferation and thus ensures functional recovery [11–13, 16]. Therefore, the representation of the body parts in the motor cortex can be altered after injury and during motor learning, and changes in the brain may be related to activity [15]. Plasticity increases because of the increased sensory input after exercises [6]. Cortical remodeling was shown to be increased after hand injuries via exercises. So, it was suggested that exercises are critical after peripheral nerve or central nervous system injuries [6]. Thus, our study looked at the effects of two different exercise programs for the upper brachial plexus lesions.

This study proved that there was no significant recovery difference between the groups in terms of time and degree, and showed that intensive exercises did not have an additional contribution on peripheral nerve injuries, which meant that the frequency of the exercises did not have a significant effect on recovery. No complications developed in the patients during their follow-ups. However, further studies may be planned with some biochemical parameters as well as cranial and spinal cord imaging, so that the effects of exercises on the recovery of cases with OBPP especially on the development of plasticity could be determined more accurately. Besides, further studies could be made on the effects of exercises on recovery with different intensity and duration as well as different frequencies.

Clinical Messages

1. Exercises positively affect recovery rate and results and prevent possible complications that may occur in cases with OBPP.

2. Intensive exercises did not have an additional contribution on peripheral nerve injuries, which means that the frequency of the exercises that the patients have daily did not have a significant effect on recovery.

Conflict of Interest: The authors declare no conflict of interest.

Financial Disclosure: The authors declared that this study has received no financial support.

REFERENCES

1. Yang LJ. Neonatal brachial plexus palsy--management and prognostic factors. *Seminars in perinatology*. 2014;38(4):222-34.
2. Chauhan SP, Blackwell SB, Ananth CV. Neonatal brachial plexus palsy: incidence, prevalence, and temporal trends. *Seminars in perinatology*. 2014;38(4):210-8.
3. Backe B, Magnussen EB, Johansen OJ, Sellaeg G, Russwurm H. Obstetric brachial plexus palsy: a birth injury not explained by the known risk factors. *Acta obstetrica et gynecologica Scandinavica*. 2008;87(10):1027-32.
4. Bahm J, Ocampo-Pavez C, Disselhorst-Klug C, Sellhaus B, Weis J. Obstetric brachial plexus palsy: treatment strategy, long-term results, and prognosis. *Deutsches Arzteblatt international*. 2009;106(6):83-90.
5. Evans-Jones G, Kay SP, Weindling AM, Cranny G, Ward A, Bradshaw A, et al. Congenital brachial palsy: incidence, causes, and outcome in the United Kingdom and Republic of Ireland. *Archives of disease in childhood Fetal and neonatal edition*. 2003;88(3):F185-9.
6. Strombeck C, Krumlind-Sundholm L, Remahl S, Sejersen T. Long-term follow-up of children with obstetric brachial plexus palsy I: functional aspects. *Developmental medicine and child neurology*. 2007;49(3):198-203.

7. Zafeiriou DI, Psychogiou K. Obstetrical brachial plexus palsy. *Pediatric neurology*. 2008;38(4):235-42.
8. Buesch FE, Schlaepfer B, de Bruin ED, Wohlrab G, Ammann-Reiffer C, Meyer-Heim A. Constraint-induced movement therapy for children with obstetric brachial plexus palsy: two single-case series. *International journal of rehabilitation research Internationale Zeitschrift für Rehabilitationsforschung Revue internationale de recherches de readaptation*. 2010;33(2):187-92.
9. Smith NC, Rowan P, Benson LJ, Ezaki M, Carter PR. Neonatal brachial plexus palsy. Outcome of absent biceps function at three months of age. *The Journal of bone and joint surgery American volume*. 2004;86-a(10):2163-70.
10. Terzis JK, Papakonstantinou KC. Management of obstetric brachial plexus palsy. *Hand clinics*. 1999;15(4):717-36.
11. Asensio-Pinilla E, Udina E, Jaramillo J, Navarro X. Electrical stimulation combined with exercise increase axonal regeneration after peripheral nerve injury. *Experimental neurology*. 2009;219(1):258-65.
12. Foret A, Quertainmont R, Botman O, Bouhy D, Amabili P, Brook G, et al. Stem cells in the adult rat spinal cord: plasticity after injury and treadmill training exercise. *Journal of neurochemistry*. 2010;112(3):762-72.
13. Kao T, Shumsky JS, Murray M, Moxon KA. Exercise induces cortical plasticity after neonatal spinal cord injury in the rat. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 2009;29(23):7549-57.
14. Bae DS, Waters PM, Zurakowski D. Correlation of pediatric outcomes data collection instrument with measures of active movement in children with brachial plexus birth palsy. *Journal of pediatric orthopedics*. 2008;28(5):584-92.
15. Strombeck C, Remahl S, Krumlinde-Sundholm L, Sejersen T. Long-term follow-up of children with obstetric brachial plexus palsy II: neurophysiological aspects. *Developmental medicine and child neurology*. 2007;49(3):204-9.
16. Molnar GE, Alexander MA. *Pediatric Rehabilitation: Hanley & Belfus*; 1999.