

The behavioral and neurochemical effects of methylprednisolone and metyrapone in a post-traumatic stress disorder rat model

 **Ayşe Melek Tanriverdi**,¹  **Banu Aydın**,¹  **Berna Terzioğlu Bebitoğlu**,²  **Hulya Cabadak**,^{1*}
 **M. Zafer Goren**^{1,3*}

¹Department of Biophysics, Marmara University Faculty of Medicine, Istanbul, Turkey

²Department of Medical Pharmacology, Istanbul Medeniyet University Faculty of Medicine, Istanbul, Turkey

³Department of Medical Pharmacology, Marmara University Faculty of Medicine, Istanbul, Turkey

ABSTRACT

OBJECTIVE: Mechanisms contributing to the post-traumatic stress disorder (PTSD) involve several physiological systems and the activation of the hypothalamic-pituitary-adrenal axis (HPA), being one of the most known system in the PTSD pathophysiology. The present study examined the effects of methylprednisolone, metyrapone and their relation with the noradrenergic system within the rostral pons a region containing the locus coeruleus (LC) in a rat model of PTSD induced with predator scent.

METHODS: Sprague-Dawley rats were exposed to the stress by exposure to the scent of dirty cat litter which is a natural stressor of a predator and 1 week later, the rats were re-exposed to a situational reminder (clean cat litter). The rats were treated either with methylprednisolone or metyrapone or physiological saline before exposure to situational reminder (n=8 in each group). Noradrenaline (NA) levels in the rostral pons homogenates were analysed by using ELISA.

RESULTS: The anxiety indices of the rats exposed to the trauma were found to be significantly higher than those of the control rats. Metyrapone produced a significant increase in the anxiety indices of the non-stressed rats and methylprednisolone did not produce a change in the anxiety indices of the non-stressed rats. Methylprednisolone treatment suppressed the anxiety in the stressed rats. Metyrapone treatment increased the anxiety indices in the stressed rats but still being lower than that of the saline treated stressed rats. Significant decrease in the freezing time was observed following the methylprednisolone treatment both in the stressed and non-stressed rats. NA content in the rostral pons of the stressed rats was significantly higher than that of the non-stressed rats. Methylprednisolone or metyrapone treatments decreased the NA content in the non-stressed rats as compared to the saline treatment. However these decreases were not significant.

CONCLUSION: The results of this study may show that stress can generate endocrine, autonomic and behavioural responses. The anxiety indices and NA levels in the rostral pons increased with the traumatic event. The methylprednisolone treatment may suppress anxiety through interactions between LC and the HPA axis.

Keywords: Anxiety index; hypothalamic-pituitary-adrenal axis; locus coeruleus; noradrenaline; rostral pons.

Cite this article as: Tanriverdi AM, Aydın B, Terzioğlu Bebitoğlu B, Cabadak H, Goren MZ. The behavioral and neurochemical effects of methylprednisolone and metyrapone in a post-traumatic stress disorder rat model. *North Clin Istanbul*

*Equally contributed to the work



Received: October 08, 2018 Accepted: May 21, 2019 Online: July 17, 2019

Correspondence: Dr. Berna TERZIOĞLU BEBITOĞLU. Marmara Üniversitesi Tıp Fakültesi, Biyofizik Anabilim Dalı, İstanbul, Turkey.
Tel: +90 532 411 49 79 e-mail: bernaterzioglu@gmail.com

© Copyright 2019 by Istanbul Provincial Directorate of Health - Available online at www.northclinist.com



This work is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License.

Post-traumatic stress disorder (PTSD) is a serious mood and anxiety problem with important manifestations for the patients, their families and the society [1]. The disease is precipitated by a traumatic event and is characterized by dysfunction in the mood lasting for more than 1 month and the persistence of signs are experiencing the stress after situational reminders, avoidance behaviours, negative mood changes and hyper-arousal or hyper-vigilance with new symptoms such as increased aggressive behaviour as stated in DSM-V [2].

Response to stress is accompanied by various emotional and physical properties as a result of activation of several physiological systems [3]. Hypothalamus-pituitary adrenal axis (HPA) activation is one of the most known responses given to stress and is accused of being involved in the PTSD pathophysiology [4, 5]. During stress, corticotropin releasing factor (CRF) acts on the pituitary gland to generate steroids that act on the monoaminergic systems involved in the stress reaction. The limbic structures such as stria terminalis and central nucleus of amygdala that are important in several autonomic and behavioural manifestations of PTSD and CRF mediates anxiety and emotional responses through these systems [6–8].

There are controversial data in the literature regarding the cortisol levels [9–13]. The change in the levels of cortisol may cause false recall of memories in the PTSD. The impact of hydrocortisone administration on automatic memory recall was investigated in a randomized, placebo-controlled study, in patients with PTSD, however hydrocortisone treatment was not found to be different than that of the placebo administration [9].

Noradrenaline (NA) is known to have a role in cognitive functions in several psychiatric disorders [14, 15]. Recently, the increased NA concentration in the rostral pons of rats demonstrated in a rat model of PTSD [16]. The Locus coeruleus (LC), a region located in the pons is rich in NA-containing neurons. NA is not only involved in the central effects occurring in the brain during stress, but it is also critical in generating the peripheral autonomic effects. It was also demonstrated that NA concentrations increase in the cerebrospinal fluid following exposure to psychological stress [17]. It was also shown that NA in the LC is usually activated together with the HPA axis activation following the stress exposure, this activation is thought to be mediated by CRF [18]. CRF projections from the central nucleus of amygdala (CeA) to the LC and the paraventricular nucleus of hypothala-

mus produce excitatory effects and cause increases in the NA input to the CeA. Cortisol may inhibit noradrenergic activity resulting in unopposed neuroendocrine response to stress [19].

Metyrapone is an agent that can prevent the synthesis of cortisol. Hence it can be used to show the interaction between the NA and the cortisol by removing the (-) negative feedback mechanism. In this setting, we wanted to test the behavioural aspects and the NA concentration in the rostral pons by using methylprednisolone or metyrapone in a rat model of PTSD induced with predator scent.

MATERIALS AND METHODS

Animals

Female Sprague–Dawley rats weighing 200–250 g were supplied from XXXX University Experimental Animal Center (XXXXXX) and local ethical committee approval (22.05.2014, 13, 2013.mar) was obtained prior to the experiments. The rats were kept in cages at $21\pm 3^{\circ}\text{C}$ and $50\pm 5\%$ humidity with reversed 12 h light/dark cycle for 10 days. The rats were fed with standard food and water ad libitum. All in vivo experimental procedures were performed in the dark phase at 10:00 am with a dim light source.

There are 6 experimental groups where each group contained 8 rats. The rats were either stressed or non-stressed and received physiological saline, metyrapone or methylprednisolone.

Predator scent test

Predator scent test was used in this study to elucidate the PTSD-like manifestations by using the scent of dirty cat litter which is a natural stressor of a predator as described previously [20]. The dirty cat litter was obtained from a male cat that used the litter for 2 days and the litter was sifted for stools. The rats are kept with in plexiglass cages (30 cm X 30 cm X 40 cm) containing 150 ml dirty cat litter for 10 min where the control (non-stressed) group was exposed to clean, unused, fresh cat litter. Following 1 week after the onset of the stress, the rats were exposed to the clean cat litter that acted as a situational reminder. The behavioural parameters on the elevated plus maze were scored from the recordings of an overhead video camera. The rats received either saline (physiological saline, PS, 0.1ml/100g) or methylprednisolone (20 mg/kg; Mustafa Nevzat, Turkey) or metyrapone

(100 mg/kg; Santa Cruz Biotechnology, Inc, CA, USA) intraperitoneal injections 10 min before the exposure to situational reminder (clean cat litter). The effective doses of methylprednisolone or metyrapone were used as stated [21, 22]. The doses were also checked in preliminary studies.

The elevated plus maze

After exposure to the situational reminder, the rats were placed on an elevated plus maze for 5 min. The elevated plus maze consists of two open (50 cm X 10 cm) and two closed (50 cm X 10 cm) arms. Closed arms were surrounded by 40 cm long walls and height of the maze was 50 cm from the ground. The labyrinth was cleaned with a 5% alcohol solution before the placement of rats.

Each rat was placed in the central square of the plus maze facing the open arms. An arm entry is defined when all four paws are present within the arm. Video recordings were used for the scoring of the entries into the open or the closed arms. The amount of time spent in the open arms provides information regarding the anxiety state of the animal. If anxiety-like feelings of animals are reduced, then the animal will spend more time in the open arms of the maze. In contrast, animals with anxiety will experience heightened anxiety-like feelings and will spend less time in the open arms of the maze. The number of entries in the open and the closed arms and the total time spent on open arms are noted and the anxiety index (N_{anxiety}) was calculated by using the following formula:

$$N_{\text{anxiety}} = 1 - \frac{1}{2} [(a/300s) + (b/c)]$$

(a=cumulative time spent in open arms (s), b=open arm entries, c=total arm entries).

The freezing behaviour is defined as the lack of all movements of rats during experiments except for respiration. The cumulative freezing time over a 5 min interval was also recorded in both stressed and non-stressed rats following exposure to the reminder. After completion of experiments, the rats were sacrificed by using high dose of pentobarbital and the selected brain regions were dissected in frozen tissues.

Preparation of the tissue samples

Upon decapitation, the rostral pons found in the brain stem was dissected. The rostral pons contains the dark coloured LC. The tissues were weighed and ground into a homogenate in the presence of ice-cold PBS (pH=7.4)

by adding 200µl per 1 mg tissue sample. After centrifugation at 3000 rpm (4°C) for 20 min, the supernatant was collected and kept at -80°C until assay. The protein content of the tissue homogenates was measured by the Lowry method [23].

Noradrenaline assay

Tissue homogenates were assayed for NA level by colorimetric method at 450 nm. NA level was detected with Rat Noradrenaline ELISA assay kit (Sunred Biotechnology Company, China), according to the manufacturers protocol. The NA content in the homogenates was adjusted according to the protein levels in the tissues and expressed as ng/mg tissue/mg protein.

Statistical analysis

All data are expressed as mean \pm SEM. The anxiety indices and NA levels of the stressed and non-stressed rats were compared with unpaired student's t-test. One-way analysis of variance and the Tukey's post hoc test were used for the analyses of anxiety indices and freezing times in all groups. For all statistical calculations, significance was accepted where $p < 0.05$. GraphPad Prism 7.0[®] is used for statistical analysis (GraphPad Software, San Diego, USA).

RESULTS

The anxiety indices of the rats exposed to the stress was measured as (0.86 \pm 0.02) and it was found to be significantly higher than those of the control rats (0.62 \pm 0.05) (Fig. 1; (unpaired student's t-test) $p < 0.05$). Metyrapone produced a significant increase in the anxiety indices of the non-stressed rats (Fig. 1; $F = 5.501$; $p = 0.0317$). Methylprednisolone did not produce a significant change in the anxiety indices of the non-stressed rats ($p = 0.9991$). The anxiety indices of the saline treated stressed rats were found to be significantly different from that of all other groups (Fig. 1b; $p < 0.0001$). Methylprednisolone treatment decreased the anxiety indices in the stressed rats (Fig. 1b; $p < 0.0001$). These findings indicate that methylprednisolone treatment suppressed anxiety. Metyrapone treatment increased the anxiety indices in the stressed rats but still being lower than that of the saline treated stressed rats ($p = 0.0001$).

The mean cumulative freezing times in all groups were analysed by using One-way analysis of variance and the treatments were found to be the source of variation both




FIGURE 1. Anxiety indices of non-stressed (**A**) and stressed rats (**B**) after injection of saline (vehicle) or methylprednisolone (20 mg/kg) or metyrapone (100 mg/kg), $p < 0.05$.




FIGURE 2. Cumulative freezing times of non-stressed (**A**) and stressed rats (**B**) after injection of saline and or methylprednisolone (20 mg/kg) or metyrapone (100mg/kg), $p < 0.05$.




FIGURE 3. Noradrenaline level after saline (a), methylprednisolone (20 mg/kg) (b) and metyrapone (100mg/kg) (c) injections in stressed and non-stressed rats, $p < 0.05$.

in stressed and nonstressed groups ($F=3.878$; $p=0.041$ and $F=21.21$; $p < 0.0001$, respectively). The post hoc test detected a significant decrease in the freezing time after methylprednisolone treatment in both stressed and non-stressed rats (Fig. 2a, b; $p=0.0358$ and $p=0.0004$, respectively). Metyrapone treatment did not produce a significant difference from the saline treatment both in the stressed and the non-stressed rats (Fig. 2, b $p=0.5715$ and $p=0.2236$, respectively).

The NA level was 7.39 ± 0.89 pg/mg in the non-stressed rats and it increased to 18.89 ± 3.37 pg/mg in the stressed rats. NA content in the rostral pons of the stressed rats was significantly higher than the that of the

non-stressed rats ($p=0.0109$). The preventive effects of drug treatments on NA levels following the exposure of the situational reminder were shown in the subsets of Figure 3. The NA concentrations of the rostral pons in the methylprednisolone treated rats were found to be 39.24 ± 10.36 pg/mg and 22.54 ± 4.78 pg/mg in the non-stressed and the stressed rats, respectively; however this decrease was not found to be statistically significant ($p=0.1688$).

NA concentration of metyrapone treated rats decreased in the stressed rats compared to the non-stressed rats being nonsignificant (77.14 ± 21.91 pg/mg vs 44.49 ± 15.29 pg/mg, respectively, $p=0.2371$).

DISCUSSION

Noradrenaline is an important neurotransmitter involved in mood, anxiety, learning, and memory functions of the central nervous system, so it has been extensively investigated in psychiatric disorders such as depression, anxiety and PTSD. The role of NA is also studied in the PTSD as the signs and symptoms imply sympathetic hyperactivity [19, 24]. NA levels in the cerebrospinal fluid were demonstrated to be increased after stress exposure (14). Previously, in the study of Terzioğlu et al. (2013), NA the levels were found to be increased in a parallel fashion with the anxiety indices in rats stressed with a predator scent [16].

We used female Sprague–Dawley rats as in the study of Aykaç et al. (2012), because the female rats were reported to develop more anxiety when compared to the male rats [25]. It was also reported that there are sex differences in stress responses and women have increased incidence of stress related psychiatric diseases [26]. Therefore we used only female rats in this study.

We investigated the effect of methylprednisolone administration on both NA level of the rostral pons and behaviours in a rat model of PTSD since glucocorticoids were shown to exert negative feedback on the HPA axis and stress causes HPA axis activation.

In PTSD studies, there are controversial data regarding the cortisol levels, some reporting the increased circulating cortisol levels, while others showing decrease [10–13]. In a study conducted on patients with chronic combat-related PTSD, higher CRF concentrations in the CSF of traumatized patients were found and researchers concluded that CRF may play role in arousal disturbances [10]. The urinary cortisol levels were also found to be below the normal range in patients with PTSD [10] and it was also noted that cortisol administration could reduce the symptoms of PTSD in chronic cases [12]. It was also shown that negative feedback inhibition of ACTH secretion by dexamethasone treatment was more apparent in PTSD patients, where no change was observed control subjects [11]. Low cortisol levels causing increased neuronal CRF transmission that activates CRF/NA circuits -between the CeA and the LC and may produce unopposed autonomic and neuroendocrine responses to stress. As cortisol causes the NA activity to decrease, it may be assumed that low levels of cortisol may make the patients to develop PTSD easily [19].

These patients may be unable to restore the increased noradrenergic activity in case of stress [19].

In our experiments, we observed that the increase in the anxiety indices of stressed rats is reversed by methylprednisolone treatment. In PTSD, the automatic recall of memories and decreased cortisol levels may be related, especially during the intrusions. However it was also demonstrated that administration of glucocorticoids following the memory reactivation results in a decrease in the subsequent recalls [27, 28].

It is well known that NA content is higher in the LC and it coordinates the structures related with physiopathology of the disease in brain regions such as the neocortex, the amygdala, the hypothalamus and the hippocampus [29, 30]. We demonstrated that noradrenergic input in the rostral pons was increased during stress. Although we detected a tendency to decrease in the NA levels in methylprednisolone or metyrapone treated stressed rats, no statistical significance was detected. The discrepancy of this finding may be due to that we measured the NA content in the rostral pons containing the LC at the end of the experiments. We do not know the time course change in the NA levels. It might have been reached to a higher level that we had detected. Microdialysis experiments will give more precise measurements.

Administration of methylprednisolone may inhibit the HPA leading to a decrease in the CRF and consequent reduction in the stimulation of the LC. This may explain the decrease in the NA in the LC following methylprednisolone administration when the stressed rats were exposed to the situational reminder. This effect was not observed with metyrapone administration, in contrast NA content was increased when compared to methylprednisolone treatment in the stressed rats. The increase in the freezing time duration was observed when the stressed rats were exposed to the reminder. The decreased freezing time may be an indicative sign of an adaptive defense response to fear exposure [31–33]. Collectively, we also observed a decrease in the freezing time in the stressed rats following methylprednisolone treatment, and this finding confirms that the corticosteroids stimulate the memory consolidation [34, 35]. The reversal of this effect with the metyrapone experiments also supports the reports regarding the association of glucocorticoids with the strong fear memories in PTSD [36]. The role of corticosteroids in the enhancement of memory consolidation during contextual fear conditioning memory was also shown in many studies [37–39].

Conclusion

Our findings also imply that glucocorticoids are pharmacologically prophylactic in stress but the inhibition of the synthesis is not effective in the stress circuitry. We concluded that the anxiety indices and NA levels in the rostral pons increased with the traumatic event. The methylprednisolone treatment may suppress anxiety through interactions between LC and the HPA axis.

Conflict of Interest: On behalf of all authors, the corresponding author states that there is no conflict of interest.

Financial Disclosure: This work was supported by Marmara University Research Fund [SAG-C-YLP-141014-0347].

Authorship Contributions: Concept – HC, MZG; Design – MZG; Supervision – HC, BA; Fundings – HC; Data collection and/or processing – AMT, BA; Analysis and/or interpretation – BTB, MZG; Writing – BTB; Critical review – MZG.

REFERENCES

- Marciniak MD, Lage MJ, Dunayevich E, Russell JM, Bowman L, Landbloom R et.al., Levine LR. The cost of treating anxiety: the medical and demographic correlates that impact total medical costs. *J Depress Anxiety* 2005;21:178-84.
- American Psychiatric Association. Diagnostic and statistical manual of mental disorders, fifth edition (DSM5), Washington, DC: American Psychiatric Association 2013.
- McEwen BS. Plasticity of the hippocampus: Adaptation to chronic stress and allostatic load. *Annals of the New York Academy of Sciences* 2001;933:265-77.
- Yehuda R. Post-traumatic stress disorder. *N Engl J Med* 2002;346:108-14.
- Yehuda R. Status of glucocorticoid alterations in post-traumatic stress disorder. *Annals of the New York Academy of Sciences* 2009;1179:56-69.
- Owens MJ, Nemeroff CB. Physiology and pharmacology of corticotropin-releasing factor. *Pharmacol Rev* 1991;43:425-73.
- Valentino RJ, Van Bockstaele E. Convergent regulation of locus coeruleus activity as an adaptive response to stress. *Eur J Pharmacol* 2008;583:194-203.
- Walker DL, Toufexis DJ, Davis M. Role of the bed nucleus of the stria terminalis versus the amygdala in fear, stress, and anxiety. *Eur J Pharmacol* 2003;463:199-216.
- Ludäscher P, Schmahl C, Feldmann RE Jr, Kleindienst N, Schneider M, Bohus M. No evidence for differential dose effects of hydrocortisone on intrusive memories in female patients with complex post-traumatic stress disorder--a randomized, double-blind, placebo-controlled, crossover study. *J Psychopharmacol* 2015;29:1077-84.
- Mason JW, Giller EL, Kosten TR, Ostroff RB, Podd L. Urinary free-cortisol levels in posttraumatic stress disorder patients. *J Nerv Ment Dis* 1986;174:145-9.
- Yehuda R, Golier JA, Halligan SL, Meaney M, Bierer LM. The ACTH response to dexamethasone in PTSD. *Am J Psychiatry* 2004;161:1397-1403.
- Yehuda R, Southwick SM, Krystal JH, Bremner D, Charney DS, Mason JW. Enhanced suppression of cortisol following dexamethasone administration in posttraumatic stress disorder. *Am J Psychiatry* 1993;150:83-6.
- Bremner JD, Licinio J, Darnell A, Krystal JH, Owens MJ, Southwick SM et.al. Elevated CSF corticotropin-releasing factor concentrations in posttraumatic stress disorder. *Am J Psychiatry* 1997;154:624-9.
- Wolf OT. The influence of stress hormones on emotional memory: relevance for psychopathology. *Acta Psychol (Amst)* 2008;127:513-31.
- Aston-Jones G, Rajkowski J, Kubiak P, Valentino RJ, Shipley MT. Role of the locus coeruleus in emotional activation. *Prog Brain Res* 1996;107:379-402.
- Terzioğlu B, Kaleli M, Aydın B, Ketenci S, Cabadak H, Gören MZ. Increased noradrenaline levels in the rostral pons can be reversed by M1 antagonist in a rat model of post-traumatic stress disorder. *Neurochem Res* 2013;38:1726-33.
- Sherin JE, Nemeroff CB. Post-traumatic stress disorder: the neurobiological impact of psychological trauma. *Dialogues Clin Neurosci* 2011;13:263-78.
- Page ME, Berridge CW, Foote SL, Valentino RJ. Corticotropin-releasing factor in the locus coeruleus mediates EEG activation associated with hypotensive stress. *Neurosci Lett* 1993;164:81-4.
- Gören MZ, Cabadak H. Noradrenaline and Post-traumatic Stress Disorder. *Comprehensive Guide to Post-Traumatic Stress Disorder*. Editors: Martin, Colin R., Preedy, Victor R., Patel, Vinood B. (Eds.) 2016.
- Pelow S, Chopin P, File SE, Briley M. Validation of open-closed arm entries in an elevated plus maze as measure of anxiety in the rat. *J Neurosci Methods* 1985;14:149-67.
- Cai WH, Blundell J, Han J, Greene RW, Powell CM. Postreactivation glucocorticoids impair recall of established fear memory. *J Neurosci* 2006;26:9560-6.
- Clay R, Hebert M, Gill G, Stapleton LA, Pridham A, Coady M et.al. Glucocorticoids are required for extinction of predator stress-induced hyperarousal. *Neurobiol Learn Mem* 2011;96:367-77.
- Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. Protein measurement with the Folin phenol reagent. *J Biol Chem* 1951;193:265-327.
- Hendrickson RC, Raskind MA. Noradrenergic dysregulation in the pathophysiology of PTSD. *Exp Neurol* 2016;284:181-95.
- Aykaç A, Aydın B, Cabadak H, Gören MZ. The change in muscarinic receptor subtypes in different brain regions of rats treated with fluoxetine or propranolol in a model of post-traumatic stress disorder. *Behav Brain Res* 2012;232:124-9.
- Bangasser DA, Valentino RJ. Sex differences in molecular and cellular substrates of stress. *Cell Mol Neurobiol* 2012;32:709-23.
- Aerni A, Traber R, Hock C, Roozendaal B, Schelling G, Papassotiropoulos A. et.al. Low-dose cortisol for symptoms of posttraumatic stress disorder. *Am J Psychiatry* 2004;161:1488-90.
- Schelling G, Kilger E, Roozendaal B, de Quervain DJ, Briegel J, Dage A. et.al. Stress doses of hydrocortisone, traumatic memories, and symptoms of posttraumatic stress disorder in patients after cardiac surgery: a randomized study. *Biol Psychiatry* 2004;55:627-33.
- Vermetten E, Bremner JD. Circuits and systems in stress. II. Applications to neurobiology and treatment in posttraumatic stress disorder. *J Depress Anxiety* 2002;16:14-38.
- Sara SJ. The locus coeruleus and noradrenergic modulation of cognition. *Nat Rev Neurosci* 2009;10:211-23.
- Knox D, Fitzpatrick CJ, George SA, Abelson JL, Liberzon I. Unconditioned freezing is enhanced in an appetitive context: implications for the contextual dependency of unconditioned fear. *Neurobiol Learn Mem* 2012;97:386-92.

32. Rosen JB. The neurobiology of conditioned and unconditioned fear: a neurobehavioral system analysis of the amygdala. *Behav Cogn Neurosci Rev* 2004;3:23-41.
33. Roelofs K. Freeze for action: neurobiological mechanisms in animal and human freezing. *Philosophical Transactions of the Royal Society of London* 2017;372:20160206.
34. Kaouane N, Porte Y, Vallée M, Brayda-Bruno L, Mons N, Calandreau L et.al. Glucocorticoids can induce PTSD-like memory impairments in mice. *Science* 2012;335:1510-3.
35. Schwabe L, Joëls M, Roozendaal B, Wolf OT, Oitzl MS. Stress effects on memory: an update and integration. *Neurosci Biobehav Rev* 2012;36:1740-9.
36. van Zuiden M, Geuze E, Willemen HL, Vermetten E, Maas M, Amarouchi K. et.al. Glucocorticoid receptor pathway components predict posttraumatic stress disorder symptom development: a prospective study. *Biol Psychiatry* 2012;71:309-16.
37. Liao Y, Shi YW, Liu QL, Zhao H. Glucocorticoid-induced enhancement of contextual fear memory consolidation in rats: Involvement of D1 receptor activity of hippocampal area CA1. *Brain Res* 2013;1524:26-33.
38. Donley MP, Schulkin J, Rosen JB. Glucocorticoid receptor antagonism in the basolateral amygdala and ventral hippocampus interferes with long-term memory of contextual fear. *Behav Brain Res* 2005;164:197-205.
39. Abrari K, Rashidy-Pour A, Semnani S, Fathollahi Y. Post-training administration of corticosterone enhances consolidation of contextual fear memory and hippocampal long-term potentiation in rats. *Neurobiol Learn Mem* 2009;91:260-5.