

Contrast-Enhanced Computed Tomography Findings of the Torsioned Wandering Spleen: A Case Report

Omer Ozcaglayan¹, Tugba Ilkem Ozcaglayan¹, Bozkurt Gulek²

ABSTRACT:

Contrast-enhanced computed tomographic findings of the wandering spleen: a case report

Objective: Wandering spleen is a rare entity that defines abnormal localization of spleen due to various causes. Wandering spleen is prone to rotate on its pedicular axis and finally torsion and infarction. Contrast enhanced computed tomography can visualize the torsioned pedicle and non enhanced parenchyma with contrast medium.

Case: A 60 year old woman who had abdominal pain was admitted to ER. Contrast enhanced computed tomography depicts the abnormal localization of spleen and absence of contrast medium in the parenchyma and pedicle. Diagnosis was torsioned wandering spleen.

Conclusion: Contrast enhanced computed tomography is very important useful modality for diagnosis of torsioned wandering spleen

Keywords: Computed tomography, torsion, wandering spleen

ÖZET:

XXXXXXXXXXXX

Amaç: Gezici dalak nadir bir antite olup, dalağın birçok sebebe bağlı olarak anormal lokalizasyonunu tanımlar. Gezici dalak, pedinküler aksı boyunca kendi eksenini etrafında dönme, sonuçta torsiyon ve enfarkt olma eğilimindedir. Kontrastlı bilgisayarlı tomografi torsiyone pedikülü ve kontrastlanmayan parankimi gösterebilir.

Olgu: Karın ağrısı olan 60 yaşında bayan hasta acil servise başvurdu. Çekilen kontrastlı bilgisayarlı tomografi dalağın anormal lokalizasyonunu, parankimin ve pedinkülün kontrastlanmadığını gösterdi. Tanı torsiyone gezici dalak olarak kondu.

Sonuç: Kontrastlı bilgisayarlı tomografi, torsiyone gezici dalak tanısında çok önemli ve yararlı bir tanı modalitesidir.

Anahtar kelimeler: Bilgisayarlı tomografi, torsiyon, gezici dalak

Ş.E.E.A.H. Tıp Bülteni XXXX;XX(x):xx-xx



¹Namık Kemal University, Faculty of Medicine, Department of Radiology, Tekirdag - Turkey
²Health Sciences University, Adana Training and Research Hospital, Department of xxxxxx, Adana - Turkey

Address reprint requests to / Yazışma Adresi:
Omer Ozcaglayan,
Namık Kemal University, Faculty of Medicine,
Department of Radiology, Tekirdag - Turkey

E-mail / E-posta:
omer.ozcaglayan@gmail.com

Date of receipt / Geliş tarihi:
July 11, 2017 / 11 Temmuz 2017

Date of acceptance / Kabul tarihi:
September 7, 2017 / 7 Eylül 2017

INTRODUCTION

The "wandering spleen" is a very rare entity, with an incidence rate estimated to be less than 0.2%. The entity has no gender predilection (1).

There are 4 ligaments which stabilize the spleen in the abdominal cavity. These are the 1) gastrosplenic, 2) colicosplenic, 3) phrenocolic, and 4) phrenosplenic, ligaments (2). A complete or incomplete embryologic developmental anomaly of these ligaments may give rise to a wandering spleen. Among the other etiological factors are

ligamentous laxity and the congenital or acquired defects of the diaphragm (3). The most important ones of the acquired factors are the advanced laxity and abdominal wall fatigue which arise due to the hormonal irregularities caused by multiparity (4). A substantially increased splenic mobility is an important cause of a torsioned wandering spleen.

It is important to remember that if there is a suspicion of torsion, imaging modalities will be more precious for early diagnosis. For diagnosis, ultrasonography and computed tomography can be used for early diagnosis.




Figure-1: Coronal oblique reformatted CECT shows the wandering spleen which is located lower abdomen, an abnormal localization for spleen (white arrows). It can clearly seen that wandering spleen is as-sociated with a torsioned pedicle (black arrow).




Figure-2: Axial CECT shows the torsioned splenic pedicle which shows 'swirling' (white arrows).

We present a female patient who was admitted to ER with abdominal pain and palpable mass lesion.

CASE REPORT

A 60-year-old female patient applied to the Emergency Department of our hospital with the complaints of nausea and vomiting during the last 10 days. The patient had undergone a right mastectomy operation due to breast cancer. Her physicals revealed a palpable midline mass in the abdomen, together with the findings of abdominal defense and rebound. Her blood pressure was 140/90 mmHg, her heart rate was 90 bpm, and her blood saturation level was measured as 92%. Her

hemogram revealed a white blood cell (WBC) count of 6700/mL (reference range being 4400 – 11000/mL), and a C reactive protein (CRP) level of 58.5 mg / L (reference range being 0 – 5 mg/L). A contrast enhanced computed tomographic (CECT) examination was performed with Toshiba Aquilion 80 Tokyo Japan. Iomeprol 300/100 mg (Iomeron 300/100 mg Bracco UK Ltd.) was used as intravenous contrast medium with 4 ml/sec injection rate. CECT disclosed that the spleen was not in place, and that there was a soft tissue lesion in the mid-abdomen. This lesion was not enhancing with iodinated contrast, and it was attached to the main vascular structures with a peduncle (Figure-1 and 2). The lesion was compatible with a wandering spleen. There was not any well known disposing factor , such as ligamental laxity or congenital malformation. It was found that the arterial peduncle feeding this wandering spleen had rotated on its axis and arterial contrast enhancing was present only at the mid section of this torsioned peduncle (Figure-3 and 4). The patient was operated and underwent an uncomplicated splenectomy procedure. Surgery revealed that the spleen was not in its normal location but was at the midline of the abdomen, and also that the peduncle was torsioned and thus the spleen necrosed.




Figure-3: In arterial phase Coronal oblique reformatted CECT image depicts that wandering spleen cannot take up contrast medium and has lost density when compared with liver. The contrast medium is discontinued in the torsioned splenic artery (black arrow).

DISCUSSION

The entity “wandering spleen” was first described by Van Horne in 1667 in an autopsy specimen (5). Hatfield et al. (6), on the other hand, were the first to define the ectopic spleen using selective angiography preoperatively, in 1976. Congenital and acquired reasons can cause wandering spleen. Congenital causes are complete or incomplete development of the ligaments of spleen. Acquired causes are increased laxity of ligaments due to hormonal changes secondary to multiparity, lymphoma and myeloproliferative disorders (7).

Patients with a wandering spleen may present




Figure-4: Sagittal reformatted image also shows the torsioned wandering spleen (white arrows). Torsioned pedicle (black arrow) and outcome of arterial contrast is well depicted (notched white arrow).

with intermittent pain attacks secondary to peduncular torsion. Cases with complete torsion may present with acute abdomen findings. Besides abdominal pain, patients may develop abdominal distension, nausea, and vomiting. Due to torsion detorsion, the pain can be present as intermittent (4). It was reported in one paper that the wandering spleen could lead to complaints of enuresis due to the ectopic spleen’s indenting pressure on the bladder. In the paper’s presented patient, the cause of the wandering spleen phenomenon was a defect in the diaphragm, and the patient’s enuresis complaints had ceased following the repair of this defect (8).

The critical question to be solved in wandering spleen cases is if there is a peduncular torsion and thus a secondary splenic ischemia and infarction.



Figure-5: Scanogram shows the intestinal structures at the spleen region (white arrows) and there is soft tissue structure associated with wandering spleen (black arrows).

This is exactly why an early diagnosis is so vital. Various imaging modalities may be used for the diagnosis. Conventional radiography, ultrasonography (US), color Doppler ultrasonography (CDUS), CT, and CECT are among these modalities (7).

Direct radiograms may show that the spleen density is missing in the normal location and there

are intestinal segments there instead. Besides, a scanogram, as was the case in our patient, may re-veal the presence of the wandering spleen as an ectopic density (Figure-5).

US, on the other hand, may not only demonstrate that the spleen is not in place, but it may also reveal the very location of the wandering spleen. US may also disclose the rotation and torsion of the splenic peduncle. With Doppler US, one can examine the vascular supply condition of both the peduncle and the splenic parenchyma. US is a cheap, easy applied modality and has no ionized radiation. But US is a user dependent modality, therefore US must be performed by experienced sonographers.

As was the case in our patient, a CECT examination may demonstrate the ectopic location of the wandering spleen, together with its increased size due to congestion secondary to torsion, and the lack of parenchymal contrast enhancement. CECT may also delineate the current status of the arterial and venous vascularization, and also the devascularized segment of the peduncle, with high accuracy. CECT has the highest contrast resolution, and therefore has the high diagnostic capability. Sagittal and coronal reformation increases the diagnostic evaluation. CT cause ionizing radiation, and this is the major disadvantage of this modality. IV contrast medium can cause side effects, such as renal impairment, anaphylaxis.

CONCLUSION

The torsion of the wandering spleen is a rare but a very important clinical entity, and it is a serious cause of acute abdomen. CECT is a very effective modality in the diagnosis of this entity, providing a fast and accurate management of the emergency situation.

REFERENCES

1. Feroci F, Miranda E, Moraldi L, Moretti R. The torsion of a wandering pelvic spleen: A case report. *Cases J* 2008; 1: 149.
2. Ostermann PA, Schreiber HW, Lierse W. The ligament system of the spleen and its significance for surgical interventions. *Langenbecks Arch Chir* 1987; 371: 207-16.
3. Cantone N, Gulia C, Miele V, Trinci M, Briganti V. Wandering Spleen and Organoaxial Gastrointestinal Volvulus after Morgagni Hernia Repair: A Case Report and Review of the Literature. *Case Rep Surgery* 2016; 2016: 1-9.
4. Yakan S, Telciler KE, Denecli AG. Acute torsion of a wandering spleen causing acute abdomen. *Hong Kong J Emerg Med* 2011; 18: 34-6.

5. Makhoulf NA, Morsy KH, Ammar S, Mohammed RA, Yousef HA, Mostafa MG. Wandering spleen in the pelvic region in an adult man with symptoms of acute abdomen. *Arab J of Gastroenterology* 2016; 17: 49-52.
6. Hatfield PM, Clouse ME, Cady B. Ectopic pelvic spleen. *Arch Surg* 1976; 111: 603-5.
7. Chauhan NS, Kumar S. Torsion of a Wandering Spleen Presenting as Acute Abdomen. *Pol J Ra-diol* 2016; 81: 110-3.
8. Emir H, Yeşildağ E, Yeker Y, Söylet Y, Şenyüz OF. A case of wandering spleen causing urinary symptoms after correction of a posterolateral diaphragmatic hernia. *Pediatr Surg Int* 2000; 16: 602-3.