



The Effects of Locally Administered Morphine Over the Dura on Postoperative Morphine Consumption and Pain After Lumbar Disc Surgery: A Prospective, Randomised, Double-Blind and Placebo-Controlled Study

Oğuzhan Kayhan , Eren Fatma Akçıl , Özlem Korkmaz Dilmen , Yusuf Tunali 

Department of Anaesthesiology and Reanimation, İstanbul University-Cerrahpaşa, Cerrahpaşa School of Medicine, İstanbul, Turkey

ORCID IDs of the authors: O.K. 0000-0001-5832-0080; E.F.A. 0000-0002-5956-2265; Ö.K.D. 0000-0002-5221-0144; Y.T. 0000-0002-6742-0128

Cite this article as: Kayhan O, Akçıl EF, Korkmaz Dilmen Ö, Tunali Y. The Effects of Locally Administered Morphine Over the Dura on Postoperative Morphine Consumption and Pain After Lumbar Disc Surgery: A Prospective, Randomised, Double-Blind and Placebo-Controlled Study. Turk J Anaesthesiol Reanim 2019; DOI: 10.5152/TJAR.2019.77854

Abstract

Objective: Effective pain management by avoiding side effects in the perioperative period is essential for patient outcome. Lumbar disc surgery is associated with moderate to severe postoperative pain, and opioids are widely used. The primary aim of the present study was to compare the effects of 1 mg and 2 mg morphine-impregnated absorbable cellulose haemostat material placed over the dura on morphine consumption, and the secondary aims were to compare pain scores and opioid-related side effects during postoperative 24 h.

Methods: The study included 44 patients (American Society of Anesthesiologists I and II). After the discectomy procedure and before the closure, in Group A (n=15), 1 mg morphine-impregnated absorbable cellulose haemostat material placed over the dura was used. In Group B (n=14), 2 mg morphine was used for the same technique, and in Group C (n=15) (control), normal saline was used. All patients used intravenous morphine patient-controlled analgesia pumps for 24 h following lumbar disc surgery. Morphine consumption, pain scores and opioid-related side effects were recorded at 10 min, 1, 2, 6, 12 and 24 h postoperatively.

Results: Morphine consumption, pain scores and opioid-related side effects were similar among the groups.

Conclusion: Morphine-impregnated absorbable cellulose haemostat material placement over the dura after single level lumbar discectomy did not reduce postoperative morphine consumption, pain scores and incidence of opioid-related side effects.

Keywords: Lumbar disc surgery, morphine, pain

Introduction

The surgical stress response due to local inflammation and systemic neuroendocrine process may affect postoperative mortality and morbidity. One of the contributing factors of surgical stress response is pain, thus pain management becomes essential in the perioperative period (1, 2).

There are many drugs and administration routes for perioperative pain management. Although systemic opioids were widely used for this purpose, their use is now limited with enhanced recovery after surgery protocols to reduce opioid-related side effects, such as sedation, respiratory depression, nausea, vomiting and ileus, in the postoperative period (3-5). Regional and local administration routes are proposed rather than systemic administration for this purpose (4, 6).

Lumbar disc surgery is associated with moderate to severe postoperative pain (7, 8). Ineffective pain management after lumbar disc surgery may cause immobilisation, thromboembolic events, pneumonia and increased sympathetic

activity leading to increased myocardial oxygen consumption (9, 10).

Thus, the primary aim of this prospective, randomised and placebo-controlled study was to compare the effects of 1 mg and 2 mg morphine-impregnated absorbable cellulose haemostat material (Surgicel®; Ethicon, NJ, USA) placed over the dura on morphine consumption during postoperative 24 h. The secondary aims were to compare pain scores and opioid-related side effects during postoperative 24 h.

Methods

This prospective, randomised, double-blind and placebo-controlled study was performed between February 2017 and April 2018. The ethics committee of Istanbul University-Cerrahpasa, Cerrahpasa School of Medicine (ethical committee no. 83045809-604.01.02; 5 January 2017) approved the study. Written informed consent was obtained from every patient. Patients with an American Society of Anesthesiologists (ASA) physical status class I–II, aged between 18 and 70 years and scheduled for microscopic single level lumbar discectomy were included in the study. Patients presenting with neurological disorders hindering communication, drug or alcohol addiction, chronic pain, allergies to any of the drugs used in the present study, hepatic or renal dysfunction and perioperative dural injury were excluded from the study.

Patients were randomised to one of three groups using a computer-generated list (in opaque sealed envelopes). All patients were previously instructed on patient-controlled analgesia (PCA) pumps (Abbott Provider®; Abbott Laboratories, Chicago, IL, USA) and visual analogue scale (VAS) from 0 to 10, with 0 being no pain and 10 being the worst pain imaginable.

Patients were sedated with intravenous (IV) midazolam (0.05 mg kg⁻¹) before surgery. In the operating room, after routine monitoring, anaesthesia was induced with propofol (2 mg kg⁻¹), rocuronium (0.5 mg kg⁻¹), remifentanyl (0.1 µg kg⁻¹) and 0.7 FiO₂ oxygen/air and maintained with sevoflurane 1 MAC in oxygen/air (FiO₂=0.40) and remifentanyl (0.05–0.1 µg kg⁻¹ h⁻¹) infusion. After orotracheal intubation, patients were then turned into the prone position supported by three surgical bolsters, one on the chest and two on the pelvis. Intraoperative analgesia was maintained with remifentanyl and tenoxicam 20 mg IV. Additional remifentanyl 25 µg IV was administered if the mean arterial pressure (MAP) and heart rate (HR) increased >20% of the baseline in all groups.

After the discectomy procedure and before the closure, in Group A (n=15), 1 mg morphine-impregnated absorbable cellulose haemostat material (Surgicel®) placed over the dura was used. In Group B (n=14), 2 mg morphine was used for

the same technique, and in Group C (n=15) (control), normal saline was used.

Sugammadex (2 mg kg⁻¹) was used to reverse residual muscle relaxation at the end of surgery. Ondansetron (8 mg IV) was administered as an antiemetic prophylaxis. All patients were extubated at the end of surgery, admitted to the post-anaesthesia care unit and then transferred to the ward when their Aldrete score reached >9 (11). All patients received IV morphine using a PCA pump for postoperative 24 h. The PCA solutions contained 100 mg morphine in 100 mL normal saline. The PCA was set to administer a bolus dose of 1 mg morphine on demand with a lockout period of 10 min and maximum 20 mg for 4 h. Two mg IV morphine was administered every 20 min in addition to PCA delivery until the pain score decreased <4.

Age, gender, ASA physical status, body mass index (BMI), duration of surgery and size of the surgical incision were recorded.

Postoperative cumulative morphine consumption and pain scores were recorded at postoperative 10 min, 1 h, 2 h, 6 h, 12 h and 24 h. Moreover, morphine-related side effects, such as nausea, vomiting, pruritus and rash, were recorded at the same time intervals and defined by a scale with 0=absent and 1=present. HR, MAPs and Ramsey sedation scores were also recorded at the same time intervals (12).

The patient and the anaesthesiologists who recorded postoperative data were blinded to the study groups.

The primary aim of the present study was to compare the effects of 1 mg and 2 mg morphine-impregnated absorbable cellulose haemostat material placed over the dura on morphine consumption during postoperative 24 h. The secondary aims were to compare pain scores and opioid-related side effects during postoperative 24 h.

Statistical analysis

A power analysis was performed prior to the study. According to the power analysis, a total of 14 patients per group should be enrolled in the study to detect at least a 10% difference in morphine consumption among the groups, with an alpha error of 0.05 and a beta error of 0.2.


Statistical analysis was performed using NCSS 10 (2015; NCSS, LLC, Kaysville, UT, USA). The Pearson chi-square test with Yates correction was used for comparison of qualitative variables between the groups, such as gender, ASA physical status and opioid-related side effects, which showed binary change. The Shapiro–Wilk normality test was used to evaluate the distribution of data. The one-way ANOVA

was used to compare normally distributed variables among the groups. The Kruskal–Wallis test was used to compare the non-normally distributed variables. The analysis of repeated measures was performed by repeated measures of ANOVA in normally distributed variables, and the Friedman test and post-hoc Wilcoxon test with Bonferroni correction were performed in non-normally distributed variables. The normally distributed values were expressed as mean (standard deviation), and the non-normally distributed values were expressed as median and interquartile range. A p value of <0.05 was considered to be statistically significant.

tion), and the non-normally distributed values were expressed as median and interquartile range. A p value of <0.05 was considered to be statistically significant.

Results

A total of 44 patients were enrolled in the study. There was no patient excluded before and after randomisation (Figure 1).



	Group A 1 mg morphine (n=15)	Group B 2 mg morphine (n=14)	Group C Control (n=15)	p
Age (year), mean (SD)	43.3 (11.0)	48.1 (13.0)	48.4 (14.7)	0.47
Gender (female/male) (n)	8/7	7/7	11/4	0.38
ASA (I/II) (n)	9/6	9/5	5/10	0.19
BMI, mean (SD)	27.0 (5.5)	28.3 (4.7)	28.5 (6.3)	0.73
Duration of surgery (min), mean (SD)	118.6 (47.1)	130 (59.9)	138.2 (51.6)	0.60
Size of incision (cm), median (IQR)	5 (5–6)	8 (5–10)	6 (5–10)	0.05*

*The size of the surgical incision was statistically longer in Group B than in Group A (p=0.05). n: number

The study groups were similar with respect to age, gender, ASA physical status scores, BMI and duration of surgery. The size of the surgical incision was statistically longer in Group B than in Group A ($p=0.05$) (Table 1).

The cumulative morphine consumption was not statistically different among the groups (Figure 2).

The groups were statistically similar with respect to VAS scores (Figure 3).

There was no statistically significant difference with respect to opioid-related side effects ($p=0.71$ for nausea and vomiting and $p=0.37$ for pruritus and rash among the groups) (Table 2).




Figure 2. Cumulative morphine consumption. The cumulative morphine consumption was not statistically different among the groups ($p>0.05$)




Figure 3. VAS scores. The groups were statistically similar with respect to VAS scores ($p>0.05$)
 VAS: visual analogue scale

The groups were similar with respect to HR, MAP and Ramsey sedation scores.

Discussion

The present study showed that locally administered morphine over the dura did not decrease postoperative morphine consumption after lumbar disc surgery. Although a statistically significant difference was not observed among the groups, the lowest morphine consumption was observed in Group A (1 mg morphine); moreover, morphine consumption was almost the same in Group B (2 mg morphine) and Group C (control). This result may be due to the longer surgical incision size in Group B (2 mg morphine). Pain scores were also lower in Group A (1 mg morphine) than in Group B (2 mg morphine) and Group C (control), but the differences were not statistically significant.

Perioperative multimodal pain management aims to provide analgesia by different neurophysiological pathways and to reduce opioid consumption, side effects and abuse. Many different modalities, such as regional and local techniques, are described for this purpose (13-16). Mastronardi et al. (17) placed 1 mg morphine-impregnated anti-adhesion gel (Adcon-L®; Gliatech Inc., Cleveland, OH, USA) over the dura after lumbar discectomy and found that analgesic requirements are lower than the control group. Wilartratsami et al. (18) placed 1 mg morphine-impregnated microfibrillar collagen sponge over the intact dural sac during single level posterior lumbar laminectomy and instrumented fusion operation and found that postoperative morphine consumption is lower than the control group during 24 h. The common feature of all of these materials is that they are composed of absorbable materials when placed in the tissue. These materials are preferred to achieve prolonged analgesic activity by the slow release of morphine. Although it is thought that the analgesia is provided by the effects of morphine on the dorsal root ganglia, a more probable mechanism of action is the opioid receptors that increase with inflammation due to surgical incision (17). The difference with these studies and our results may be due to our material properties and the ability of morphine absorbance. The material that we use in our institution may not be appropriate for this purpose. We recommend the use of gel or sponge materials for future studies.

Table 2. Opioid-related side effects

	Group A 1 mg morphine (n=15)	Group B 2 mg morphine (n=14)	Group C Control (n=15)	p
Nausea–vomiting (n)	6	6	4	0.71
Pruritus–rash (n)	3	0	0	0.37

The groups were statistically similar with respect to opioid-related side effects ($p>0.05$). n: number

Wu et al. (4) placed 1 mg morphine-impregnated microfibrillar haemostasis sponge (Avitene®; Davol Inc., Cranston, RI, USA) on the dura at the end of the posterior lumbar spinal decompression and fusion operations and reported that pain scores are similar with IV morphine PCA and are statistically lower than intramuscular meperidine. Chen et al. (19) found that 3 mg morphine and 80 mg methylprednisolone-impregnated microfibrillar collagen placed over the dural sac improve pain scores after lumbar multilevel laminectomy. Mastronardi et al. (17) found that pain scores are statistically lower in the 1 mg morphine-impregnated anti-adhesion gel (Adcon-L®) group than in the control group in lumbar discectomy. Wilartratsami et al. (18) also found that pain scores are lower in the 1 mg morphine-impregnated microfibrillar collagen sponge group than in the control group. We did not observe a statistically significant difference with respect to pain scores among the groups. Pain scores were <5 in all groups. Spinal fusion surgery is one of the most painful surgeries (20). Different results in pain scores may be due to the difference in the types of surgery. Our study was single level lumbar discectomy, whereas other previous studies were spinal decompression with instrumented fusion operations where pain intensity could be more severe.

The systemic administration of opioids even with PCA is associated with dose-dependent side effects, such as sedation, respiratory depression, nausea, vomiting and ileus (21). It is demonstrated that a low dose of epidural morphine administration (10–20 µg kg⁻¹) provided adequate analgesia, but vomiting was not reduced even with low dose (22). We evaluated two doses of morphine in our study and did not find any difference with respect to side effects. Wu et al. (4) found postoperative nausea and vomiting to be lower in the morphine-impregnated microfibrillar haemostasis sponge (Avitene®) group than in the IV morphine PCA group. We administered prophylactic antiemetic ondansetron intraoperatively, and they used IV morphine PCA continuously; therefore, postoperative nausea and vomiting were lower in our study than in Wu et al.'s study (4).

Our study groups were similar with regard to age, gender, ASA physical status scores, duration of surgery that may have influence on postoperative pain scores and development of opioid-related side effects (23-28).

The present study has some limitations. First, it is powered to determine a 10% difference in morphine consumption; perhaps a bigger sample size was needed to analyse the difference in pain scores. Second, remifentanyl as analgesic was used intraoperatively. Remifentanyl is supposed to have an opioid-induced hyperalgesia (OIH) effect and may cause acute opioid tolerance that may confound our results (29). The activation of spinal N-methyl-D-aspartate receptors by pure µ receptor

agonists, such as remifentanyl, is proposed as the main mechanism of OIH (30). Intraoperative remifentanyl infusion was found to be associated with increased early (within postoperative 4 h) postoperative pain and morphine requirements (31). The higher pain scores in the early postoperative period in our study might be due to this effect.

Conclusion

Morphine-impregnated absorbable cellulose haemostat material (Surgicel®) placement over the dura did not reduce postoperative morphine consumption, pain scores and incidence of opioid-related side effects; more larger sample-sized studies are needed.

Ethics Committee Approval: Ethics committee approval was received for this study from the ethics committee of Istanbul University-Cerrahpasa, Cerrahpasa School of Medicine (83045809-604.01.02; 5 January 2017).

Informed Consent: Written informed consent was obtained from patients who participated in this study.

Peer-review: Externally peer-reviewed.

Author Contributions: Concept – O.K., Y.T. Design – O.K., Y.T.; Supervision – Ö.K.D., Y.T. ; Resources – Ö.K.D., Y.T. Materials – O.K.; Data Collection and/or Processing – O.K. ; Analysis and/or Interpretation – O.K., E.F.A., Ö.K.D., Y.T.; Literature Search – O.K.; Writing Manuscript – O.K., Ö.K.D., Y.T., E.F.A.; Critical Review – O.K.D., Y.T.

Conflict of Interest: The authors have no conflicts of interest to declare.

Financial Disclosure: The authors declared that this study has received no financial support.

References

1. Lindley EM, Milligan K, Farmer R, Burger EL, Patel VV. Patient-controlled transdermal fentanyl versus intravenous morphine pump after spine surgery. *Orthopedics* 2015; 38: 819-24. [\[CrossRef\]](#)
2. Hurley RW, Murphy JD, Wu CL. Acute postoperative pain. In: Miller RD, editor. *Miller's Anesthesia*. 8th ed. Philadelphia: Elsevier/Saunders; 2015. pp.2974-98.
3. Polat R, Peker K, Gülöksüz ÇT, Ergil J, Akkaya T. Comparison of the postoperative analgesic effects of paracetamol-codeine phosphate and naproxen sodium-codeine phosphate for lumbar disk surgery. *Kaohsiung J Med Sci* 2015; 31: 468-72. [\[CrossRef\]](#)
4. Wu MH, Wong CH, Niu CC, Tsai TT, Chen LH, Chen WJ. A comparison of three types of postoperative pain control after posterior lumbar spinal surgery. *Spine* 2011; 36: 2224-31. [\[CrossRef\]](#)
5. Kleppe KL, Greenberg JA. Enhanced Recovery After Surgery Protocols: Rationale and Components. *Surg Clin North Am* 2018; 98: 499-509. [\[CrossRef\]](#)

6. Brandal D, Keller MS, Lee C, Grogan T, Fujimoto Y, Gricourt Y, et al. Impact of Enhanced Recovery After Surgery and Opioid-Free Anesthesia on Opioid Prescriptions at Discharge From the Hospital: A Historical-Pro prospective Study. *Anesth Analg* 2017; 125: 1784-92. [\[CrossRef\]](#)
7. Korkmaz Dilmen O, Tunali Y, Cakmakkaya OS, Yentur E, Tutuncu AC, Tureci E, et al. Efficacy of intravenous paracetamol, metamizol and lornoxicam on postoperative pain and morphine consumption after lumbar disc surgery. *Eur J Anaesthesiol* 2010; 27: 428-32. [\[CrossRef\]](#)
8. Tunali Y, Akcil EF, Dilmen OK, Tutuncu AC, Koksak GM, Akbas S, et al. Efficacy of intravenous paracetamol and dexketoprofen on postoperative pain and morphine consumption after a lumbar disc surgery. *J Neurosurg Anesthesiol* 2013; 25: 143-7. [\[CrossRef\]](#)
9. Macres SM, Moore PG, Fishman SM. Acute pain management. In: Barash PG, Cullen BF, Bruce F, Stoelting RK, Cahalan MK, Stock MC, Ortega R, Sharar SM, Holt NF editors. *Clinical Anesthesia*. 8th ed: Philadelphia: Lippincott Williams & Wilkins; 2017; 1562-606.
10. Kehlet H, Holte K. Effect of postoperative analgesia on surgical outcome. *Br J Anaesth* 2001; 87: 62-72. [\[CrossRef\]](#)
11. Camerani S, Capuzzo M, Ieffa E, Pescolderung M, Braccini L, Volta CA. Total intravenous anesthesia with superficial cervical block or morphine transition in patients undergoing carotid endarterectomy. *Minerva Anesthesiol* 2014; 80: 676-84.
12. Reschreiter H, Maiden M, Kapila A. Sedation practice in the intensive care unit: a UK national survey. *Crit Care* 2008; 12: 152. [\[CrossRef\]](#)
13. Beverly A, Kaye AD, Ljungqvist O, Urman RD. Essential elements of multimodal analgesia in enhanced recovery after surgery (ERAS) guidelines. *Anesthesiol Clin* 2017; 35: 115-43. [\[CrossRef\]](#)
14. Kehlet H. Multimodal approach to control postoperative pathophysiology and rehabilitation. *Br J Anaesth* 1997; 78: 606-17. [\[CrossRef\]](#)
15. Feldheiser A, Aziz O, Baldini G, Cox BP, Fearon KC, Feldman LS, et al. Enhanced recovery after surgery (ERAS) for gastrointestinal surgery, part 2: consensus statement for anaesthesia practice. *Acta Anaesthesiol Scand* 2016; 60: 289-334. [\[CrossRef\]](#)
16. Wick EC, Grant MC, Wu CL. Postoperative multimodal analgesia Pain management with nonopioid analgesics and techniques: a review. *JAMA Surg* 2017; 152: 691-7. [\[CrossRef\]](#)
17. Mastronardi L, Pappagallo M, Puzzilli F, Tatta C. Efficacy of the morphine-Adcon-L compound in the management of postoperative pain after lumbar microdiscectomy. *Neurosurgery* 2002; 50: 518-25. [\[CrossRef\]](#)
18. Wilartratsami S, Sanansilp V, Ariyawatkul T, Chotigavanichaya C, Pichaisak W, Chotivichit A, et al. The effect of epidural low-dose morphine-soaked microfibrillar collagen sponge in postoperative pain control after laminectomy and instrumented fusion: a randomized double-blind placebo-controlled study. *J Med Assoc Thai* 2014; 97(Suppl 9): 62-7.
19. Chen TY. Efficacy of topical microfibrillar collagen mixed with steroid hormone and morphine for postoperative pain control during lumbar laminectomy: a preliminary report. *Chang Gung Med J* 2002; 25: 81-8.
20. Gerbershagen HJ, Aduckathil S, van Wijck AJ, Peelen LM, Kalkman CJ, Meissner W. Pain intensity on the first day after surgery: a prospective cohort study comparing 179 surgical procedures. *Anesthesiology* 2013; 118: 934-44. [\[CrossRef\]](#)
21. Cohen BE, Hartman MB, Wade JT, Miller JS, Gilbert R, Chapman TM. Postoperative pain control after lumbar spine fusion. Patient-controlled analgesia versus continuous epidural analgesia. *Spine* 1997; 22: 1892-6. [\[CrossRef\]](#)
22. Castillo-Zamora C, Castillo-Peralta LA, Nava-Ocampo AA. Dose minimization study of single-dose epidural morphine in patients undergoing hip surgery under regional anesthesia with bupivacaine. *Paediatr Anaesth* 2005; 15: 29-36. [\[CrossRef\]](#)
23. Taylor S, Kirton OC, Staff I, Kozol RA. Postoperative day one: a high risk period for respiratory events. *Am J Surg* 2005; 190: 752-6. [\[CrossRef\]](#)
24. Sarton E, Teppema L, Dahan A. Sex differences in morphine-induced ventilatory depression reside within the peripheral chemoreflex loop. *Anesthesiology* 1999; 90: 1329-38. [\[CrossRef\]](#)
25. Gupta K, Prasad A, Nagappa M, Wong J, Abrahamyan L, Chung FF. Risk factors for opioid-induced respiratory depression and failure to rescue: a review. *Curr Opin Anaesthesiol* 2018; 31: 110-9.
26. Mordhorst C, Latz B, Kerz T, Wisser G, Schmidt A, Schneider A, et al. Prospective assessment of postoperative pain after craniotomy. *J Neurosurg Anesthesiol* 2010; 22: 202-6. [\[CrossRef\]](#)
27. Celentano DD, Linet MS, Stewart WF. Gender differences in the experience of headache. *Soc Sci Med* 1990; 30: 1289-95. [\[CrossRef\]](#)
28. Notermans SL, Tophoff MM. Sex difference in pain tolerance and pain apperception. *Psychiatr Neurol Neurochir* 1967; 70: 23-9.
29. Santonocito C, Noto A, Crimi C, Sanfilippo F. Remifentanyl-induced postoperative hyperalgesia: current perspectives on mechanisms and therapeutic strategies. *Local Reg Anesth* 2018; 11: 15-23. [\[CrossRef\]](#)
30. Simonnet G. Opioids: from analgesia to anti-hyperalgesia? *Pain* 2005; 118: 8-9. [\[CrossRef\]](#)