



# Diagnostic Challenge of an Ulnar Nerve Schwannoma Confused with a Lipoma

## *Lipom ile Karışan Ulnar Sinir Schwannomu*

Suna Örs<sup>1</sup>, Atilla Kokurcan<sup>1</sup>, Yasemin Kayadibi<sup>1</sup>, Turgut Kayadibi<sup>2</sup>

<sup>1</sup>Van Area Training and Research Hospital, Clinic of Radiology, Van, Turkey

<sup>2</sup>Van Area Training and Research Hospital, Clinic of Plastic Surgery, Van, Turkey

**Keywords:** Lipoma, schwannoma, paresthesia, ulnar nerve

**Anahtar Kelimeler:** Lipom, schwannom, parestezi, ulnar sinir

Dear Editor,

Schwannomas are benign tumors that originate from Schwann cells. These benign tumors are easily mistaken for other entities such as lipomas, neurofibromas, hemangiomas, and synovial cysts (1,2). Here, we report a case of schwannoma of the ulnar nerve that was mistaken for a lipoma in a patient who had a history of lipoma, as surgically diagnosed. The purpose of this report was to emphasize the importance of appropriate diagnosis and management of peripheral sheath nerve tumors and highlight the role of imaging features in the suspicion of schwannomas.

A man aged 59 years presented with a five-year history of a slowly enlarging mass on his right distal forearm. The patient had a history of similar lesions on the right and left upper limbs that were diagnosed as lipoma through surgery. During the last six months, the mass had been associated with dull pain and discomfort. His initial neurologic examination revealed a minimally mobile 2x1-cm lesion on the ulnar aspect of the forearm associated with dull pain triggered by mild pressure. Hoffman-Tinel's sign was positive and there was minimal paresthesia over the 4<sup>th</sup> and 5<sup>th</sup> fingers of the right hand. An ultrasonography examination revealed an 8x15-mm, fusiform solid mass near the ulnar artery on the right distal forearm. The non-vascular mass was well-defined, homogeneous, isoechoic with the ulnar nerve, and had continuity with the ulnar nerve

(Figure 1A, 1D). Magnetic resonance imaging revealed that the tumor was isointense to surrounding muscles on T1-weighted images, and hyperintense on T2-weighted images. Homogeneous contrast enhancement was observed (Figure 1B, 1E). A histopathologic examination established the diagnosis of classic schwannoma (Figure 1C, 1F).

Clinically, schwannomas are slow-growing tumors, sometimes associated with pain and paresthesia (3).

Ulnar nerve schwannoma cases have rarely been reported in the literature (4). Careful physical examination and imaging features raise the diagnostic suspicion of schwannomas, and histopathologic features are the mainstay for the definitive diagnosis of these tumors (5). When a clinician encounters a surface lesion that appears benign, peripheral nerve sheath tumors should be considered as a possible differential diagnosis.

### Ethics

**Informed Consent:** Consent form was filled out by all participants.

**Peer-review:** Internally peer-reviewed.

### Authorship Contributions


Surgical and Medical Practices: T.K., Concept: S.Ö., Design: S.Ö., Data Collection or Processing: A.K., Analysis or Interpretation: Y.K., Literature Search: Y.K., Writing: S.Ö.

**Address for Correspondence/Yazışma Adresi:** Yasemin Kayadibi MD, Van Area Training and Research Hospital, Clinic of Radiology, Van, Turkey

Phone: +90 536 257 08 79 E-mail: ysmnkayadibi@gmail.com ORCID ID: orcid.org/0000-0003-1590-5382

**Received/Geliş Tarihi:** 24.11.2016 **Accepted/Kabul Tarihi:** 31.01.2017

©Copyright 2017 by Turkish Neurological Society  
Turkish Journal of Neurology published by Galenos Publishing House.



**Figure 1.** The round-shaped homogenous, isoechoic, non-vascular mass (arrow) was observed on color Doppler image (A). Homogenous contrast enhancement of the lesion (arrow) on axial T1-weighted images after contrast administration (B). Histopathologic image of the schwannoma, (hematoxylin&eosin stain) (C). The non-vascular mass (arrow) had continuity with and was isoechoic with the ulnar nerve (D). The normal ulnar nerve of the opposite side on a T1-weighted image (E). Histopathologic image of the schwannoma; common nuclear and cytoplasmic positivity for S-100 (F)

**Conflict of Interest:** No conflict of interest was declared by the authors.

**Financial Disclosure:** The authors declared that this study received no financial support.

## References

1. Morris JH. The nervous system. In: Cotran RS, Kumar V, Robbins SL (eds). Robbins Pathologic Basis of Disease. 4th ed. Philadelphia: WB Saunders, 1989:1445-1447.
2. Gupta G, Maniker A. Malignant peripheral nerve sheath tumours. *Neurosurg Focus* 2007;22:12.
3. Komurcu E, Kaymaz B, Adam G, Gokmen F, Murath A. A hybrid schwannoma of median nerve. *J Coll Physicians Surg Pak* 2014;24(Suppl 1):32-33.
4. Vigler M, Levine LJ, Posner MA. Multiple neurilemmomas in the upper extremity: a series of three cases. *Bull NYU Hosp Jt Dis* 2008;66:61-64.
5. Adani R, Baccarani A, Guidi E, Tarallo L. Schwannomas of the upper extremity: diagnosis and treatment. *Chir Organi Mov* 2008;92:85-88.