



Shift in Time Zone: The Biologic Clock and Neurology

Zaman Dilimi Değişikliği: Biyolojik Saat ve Nöroloji

● Feray Güleç¹, ● Raika Durusoy², ● Neşe Çelebisoy³

¹Izmir Tepecik Training and Research Hospital, Clinic of Neurology, Izmir, Turkey

²Ege University Faculty of Medicine, Department of Public Health, Izmir, Turkey

³Ege University Faculty of Medicine, Department of Neurology, Izmir, Turkey

Keywords: Shift in time zone, daylight saving, biological clock

Anahtar Kelimeler: Zaman dilimi değişikliği, yaz saati, biyolojik saat

Introduction

Almost every tissue in the human body functions according to an internal clock regulated to a 24-hour circadian rhythm. Tissues and cells that begin to function outside of this rhythm have a tendency toward carcinogenesis (1). The hypothesis of the mechanism of the biologic clock has brought a Nobel prize to its owner (2). With this communication, we emphasize the relationship between the biologic clock and neurology and aim to inspire readers in challenging fields of research.

The 'summer-time' implementation has been rendered permanent in Turkey according to an enactment published in the Official Gazette (No: 29825) on September 8th, 2016, which entered into force the same day. As we do not in fact have a real 'winter time', this can be considered as a 'shift in time zone.' The break-up of the bond between sleep and its most powerful 'zeitgeber' daylight can be searched with the term 'daylight saving time' in the medical literature, and it can be seen that this situation has been the subject of many studies varying from myocardial infarction to the occurrence of stroke (3).

The suprachiasmatic nucleus located in the hypothalamus plays an important role in the regulation of the circadian rhythm. The biologic clock relating to different anatomic structures and neuroendocrine processes has a principal role in the secretion of melatonin, variations in the temperature of the body, cortisol release, and appetite. It also acts in modalities such as attention/concentration, mood, and metabolic and autonomic modulation. The

consequences of sleep disorders due to external causes incompatible with the biorhythm can be much more than just physical tiredness. This situation influences perception, data processing, concentration, and recalling functions of individuals in workplaces, in traffic, and at school or home. Waking up with daylight at the last part of a normal night sleep affects daily mental performance positively and facilitates falling asleep at night, and enables preparation for the next day. The response to suppress the effects of environmental time determinants has negative impacts both on cognitive performance and the physical "sense of well-being". Young people and children in particular can encounter problems, which might impact the following years of their lives (4).

Not wearing a biologic clock affects the hypothalamopituitary axis (HPA) with a potential to result in unfavorable outcomes such as obesity, diabetes, and hypertension (5). Epileptic seizures, migraine episodes, and mood disorders can be exacerbated. Stroke and cardiovascular events such as myocardial infarction might develop (4). Impacts on the HPA working in close contact with the biologic clock might alter the attack frequency and severity of autoimmune diseases. The implementation of a 'shift in time zone' enables a unique opportunity to scientists in Turkey to design epidemiologic studies to better understand the link between the biologic clock/circadian rhythm and stroke, migraine, epilepsy, episodes of autoimmune diseases such as multiple sclerosis, myasthenia gravis, and of course sleep disorders. We hope that this paper inspires our colleagues to design studies leading to interesting scientific results.

Address for Correspondence/Yazışma Adresi: Feray Güleç MD, Izmir Tepecik Training and Research Hospital, Clinic of Neurology, Izmir, Turkey

Phone: +90 505 718 81 98 E-mail: drferaygulec@gmail.com ORCID ID: orcid.org/0000-0001-7113-1145

Received/Geliş Tarihi: 08.02.2018 **Accepted/Kabul Tarihi:** 05.04.2018

©Copyright 2018 by Turkish Neurological Society

Turkish Journal of Neurology published by Galenos Publishing House.

Ethics

Peer-review: Internally peer-reviewed.

Authorship Contributions

Concept: F.G., R.D., N.Ç., Design: F.G., R.D., N.Ç., Data Collection or Processing: F.G., R.D., N.Ç., Analysis or Interpretation: F.G., R.D., N.Ç., Literature Search: F.G., R.D., N.Ç., Writing: F.G., R.D., N.Ç.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study received no financial support.

References

1. Sancar A, Lindsey-Boltz LA, Gaddameedhi S, et al. Circadian Clock, Cancer, and Chemotherapy. *Biochemistry* 2015;54:110-123.
2. https://www.nobelprize.org/nobel_prizes/medicine/laureates/2017/press.html
3. Sipilä JO, Ruuskanen JO, Rautava P, Kytö V. Changes in ischemic stroke occurrence following daylight saving time transitions. *Sleep Med* 2016;27:20-24.
4. Boergers J, Gable CJ, Owens JA. Later school start time is associated with improved sleep and daytime functioning in adolescents. *J Dev Behav Pediatr* 2014;35:11-17.
5. Parsons MJ, Moffitt TE, Gregory AM, et al. Social jetlag, obesity and metabolic disorder: investigation in a cohort study. *Int J Obes (Lond)* 2015;39:842-848.