



Inflammation with Neutrophil-to-lymphocyte Ratio and Platelet-to-lymphocyte Ratio in Restless Legs Syndrome

Huzursuz Bacaklar Sendromunda Enflamasyonun Nötrofil-lenfosit ve Trombosit-lenfosit Oranları ile Değerlendirilmesi

Ali Zeynal Abidin Tak¹, Yıldızhan Şengül²

¹Adiyaman University Faculty of Medicine, Department of Neurology, Adiyaman, Turkey

²Bezmialem Vakif University Faculty of Medicine, Department of Neurology, Istanbul, Turkey

Keywords: Restless legs syndrome, neutrophil-to-lymphocyte ratio, platelet-to-lymphocyte ratio

Anahtar Kelimeler: Huzursuz bacaklar sendromu, nötrofil-lenfosit oranı, platelet-lenfosit oranı

Dear Editor,

Restless legs syndrome (RLS) is a chronic, progressive sensorimotor neurologic condition (1). In some recent studies, RLS has been associated with inflammatory diseases such as rheumatoid arthritis, systemic lupus, Crohn's disease, and celiac disease (2,3), and the possible role of inflammation in the etiology has been discussed. Hypoxia-inducible factor-1 alpha (HIF-1a) acts as a regulator in inflammation (4). In a study by Patton et al. (5), high HIF-1a values were shown in the substantia nigra neurons of patients with RLS, indicating the role of inflammation at the cellular level in RLS. The neutrophil-to-lymphocyte ratio (NLR) and platelet-to-lymphocyte ratio (PLR) are low cost and easily applicable methods that have recently been studied as markers of inflammation in various studies. NLR has been evaluated in a number of central and systemic neurologic diseases such as ischemic and hemorrhagic cerebrovascular diseases, myasthenia gravis, and multiple sclerosis, and its association with prognosis has been shown (2,3). In the literature, a single study evaluated inflammation with NLR, which was found to be significantly higher in the RLS group and the effect of inflammation in the etiology was discussed (3). In our study, we aimed to evaluate the inflammation with NLR and PLR in RLS and to discuss whether they may be biomarkers in terms of inflammation in such neurological diseases.

Ethics

Informed Consent: Consent form was filled out by all participants.

Peer-review: Internally peer-reviewed.

Authorship Contributions

Surgical and Medical Practices: A.Z.A.T., Concept: A.Z.A.T., Design: Y.Ş., Data Collection or Processing: A.Z.A.T., Analysis or Interpretation: Y.Ş., Literature Search: A.Z.A.T., Writing: A.Z.A.T.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study received no financial support.

References

- Allen RP, Picchietti DL, Garcia-Borreguero D, et al. International Restless Legs Syndrome Study Group. Restless legs syndrome/Willis-Ekbom disease diagnostic criteria: updated International Restless Legs Syndrome Study Group (IRLSSG) consensus criteria--history, rationale, description, and significance. *Sleep Med* 2014;15:860-873.
- Tak AZA, Sengül Y. Evaluation of neutrophil-to-lymphocyte ratio and platelet-to-lymphocyte ratio in essential tremor. *Idoggyogy Sz* 2019;72:33-38.
- Varım C, Acar BA, Uyanık MŞ, et al. Association between the neutrophil-to-lymphocyte ratio, a new marker of systemic inflammation, and restless legs syndrome. *Singapore Med J* 2016;57:514-516.
- Imtiyaz HZ, Simon MC. Hypoxia-inducible factors as essential regulators of inflammation. *Curr Top Microbiol Immunol* 2010;345:105-120.
- Patton SM, Ponnuru P, Snyder AM, Podskalny GD, Connor JR. Hypoxia-inducible factor pathway activation in restless legs syndrome patients. *Eur J Neurol* 2011;18:1329-1335.

Address for Correspondence/Yazışma Adresi: Ali Zeynal Abidin Tak MD, Adiyaman University Faculty of Medicine, Department of Neurology, Adiyaman, Turkey

Phone: +90 530 606 19 32 E-mail: atak@adiyaman.edu.tr ORCID: orcid.org/0000-0002-3783-184X

Received/Geliş Tarihi: 01.01.2019 **Accepted/Kabul Tarihi:** 03.03.2019

©Copyright 2019 by Turkish Neurological Society
Turkish Journal of Neurology published by Galenos Publishing House.