

# Comparison of the demographic and clinical features of pregnant and non-pregnant patients undergoing appendectomy

✉ Cemalettin Koç, M.D.,<sup>1</sup> ✉ Sami Akbulut, M.D.,<sup>1</sup> ✉ Ebru İnci Coşkun, M.D.,<sup>2</sup>  
✉ Barış Sarıcı, M.D.,<sup>1</sup> ✉ Sezai Yılmaz, M.D.<sup>1</sup>

<sup>1</sup>Department of Surgery and Liver Transplant Institute, İnönü University Faculty of Medicine, Malatya-Turkey

<sup>2</sup>Department of Obstetrics and Gynecology, İnönü University Faculty of Medicine, Malatya-Turkey

## ABSTRACT

**BACKGROUND:** This retrospective study aims to compare the demographic and clinicopathological characteristics of the pregnant and non-pregnant patients who underwent appendectomy with a presumed diagnosis of acute appendicitis.

**METHODS:** Between June 2009 and January 2019, 431 reproductive-aged of female patients underwent appendectomy with a presumed diagnosis of acute appendicitis. Patients were divided into two groups considering their pregnancy status: pregnant group (n=48) and non-pregnant group (n=383). Both groups were compared with respect to demographic, clinical and histopathological features.

**RESULTS:** No statistically significant difference was found between pregnant and non-pregnant groups except total bilirubin level (p=0.019) and ultrasonographic findings (p=0.016). In the non-pregnant group, negative appendectomy and perforation rates were 26% and 10.5%, where these rates for the pregnant group were 20.8% and 4.2%. Sensitivity, specificity and accuracy rates of ultrasonography for the pregnant group were 50%, 100% and 58.5%, where these rates for the non-pregnant group were 67.3%, 57.9% and 65%. The pregnancy date was the first trimester in 52.1%, the second trimester in 29.2% and the third trimester in 16.7% of the pregnant patients. None of the term births (87.5%) resulted in neither a fetal nor a maternal complication. However, 12.5% of the preterm births resulted in neonatal mortality.

**CONCLUSION:** Although not statistically significant, this study points out relatively lower rates of negative appendectomy and perforated acute appendicitis among pregnant patients, which is related to the overly attentive evaluation of pregnant patients admitted due to acute abdomen.

**Keywords:** Acute appendicitis; appendectomy; obstetric complications; pregnancy; preterm labour.

## INTRODUCTION

Acute appendicitis (AAP) is among the leading causes of emergency unit admissions due to abdominal pain and appendectomy is among the world wide most performed surgical procedures.<sup>[1,2]</sup> Obstruction of appendix vermiformis lumen due to any cause triggers an inflammatory process that initially begins in epithelium progressing into serosa, resulting in classical sign and symptoms of AAP.<sup>[2]</sup> Lifetime real AAP risk varies from 5% to 20% and which is around 6.9% for women.<sup>[1,2]</sup>

AAP is one of the most frequent conditions of pregnant women requiring an emergent surgical procedure. AAP incidence during pregnancy varies from 1/800 to 1/1500 and has a relatively lower incidence compared with non-pregnant women of the same age.<sup>[3]</sup> Loss of appetite, nausea, and vomiting, abdominal pain are the cardinal signs and symptoms of AAP, which are frequently common in the normal physiological course of pregnancy.<sup>[4]</sup> Therefore, diagnosing AAP in a pregnant patient is challenging and 25% to 50% of patients are preoperatively underdiagnosed.<sup>[4]</sup> Delay in diagnosis or

Cite this article as: Koç C, Akbulut S, Coşkun Eİ, Sarıcı B, Yılmaz S. Comparison of the demographic and clinical features of pregnant and non-pregnant patients undergoing appendectomy. *Ulus Travma Acil Cerrahi Derg* 2020;26:555-562.

Address for correspondence: Cemalettin Koç, M.D.

İnönü Üniversitesi Tıp Fakültesi, Genel Cerrahi Anabilim Dalı, Malatya, Turkey

Tel: +90 422 - 341 06 60 E-mail: ckoc\_41@hotmail.com

*Ulus Travma Acil Cerrahi Derg* 2020;26(4):555-562 DOI: 10.14744/tjtes.2020.12544 Submitted: 19.09.2019 Accepted: 15.01.2020 Online: 15.06.2020  
Copyright 2020 Turkish Association of Trauma and Emergency Surgery



underdiagnosis in pregnant patients results in perforation and peritonitis, which leads to unfavorable complications of early delivery, miscarriage, fetal loss, and maternal mortality.<sup>[4]</sup> This study presented aims, first of all, compare demographic, biochemical and histopathological features of pregnant and non-pregnant female patients admitted to our surgery clinic instruction with an initial diagnosis of AAP. The second aim is to present maternal and fetal complication following appendectomy among pregnant patients.

## MATERIALS AND METHODS

Between June 2009 and January 2019, the demographic, biochemical and histopathological features of the 48 pregnant patients who underwent appendectomy with presumed diagnosis of AAP at Inonu University Faculty of Medicine, Department of Surgery were analyzed retrospectively. This group was defined as the Pregnant group (n=48). A control group was created to compare with the pregnant group and this group was defined as non-pregnant group (n=383). The non-pregnant group consists of reproductive-aged (range: 18–45 years) female patients who presented to our emergency unit with abdominal pain at the same time frame and underwent appendectomy with the presumed diagnosis of AAP. Patients' medical records were reviewed after obtaining approval from Inonu University institutional review board for non-interventional studies (Approval No:2019/4-41). Both groups were compared in terms of age (years), white blood cell (WBC), Neutrophil, Lymphocyte, Platelets, mean corpuscular hemoglobin (MCH), red cell distribution width (RDW), mean platelet volume (MPV), mean corpuscular volume (MCV), platelet distribution width (PDW), C-reactive protein (CRP), neutrophil to lymphocyte ratio (NLR), platelet to lymphocyte ratio (PLR), platelet to neutrophil ratio (PNR), white blood cell to lymphocyte ratio (WLR), white blood cell to neutrophil ratio (WNR) bilirubin, appendix width (mm), appendix length (mm) and histopathological findings.

Pregnancy is categorized into three phases of the first trimester (0–14 wk), second trimester (15–28 wk) and third trimester (29–42 wk).<sup>[5]</sup> Patients younger than 18 years of age are excluded from this study. Therefore, non-pregnant patients from 18 to 45 years of age are included in the control group for matching the age of the pregnant group. All patients admitted to the emergency unit with an initial diagnosis of AAP were evaluated with routine anamnesis (such as date of last menstrual period, sexual activity), blood  $\beta$ -hCG level and abdominopelvic ultrasonography (US) to rule out obstetrical and gynecological conditions.

Demonstration of a non-compressible, aperistaltic tubular structure originating from cecum, with a blind end, anteroposterior diameter >6 mm in US evaluation is defined as AAP. Additionally, thickened intestinal wall, inflammation, increased echogenicity of surrounding mesenteric fatty tissue, heterogeneity, appendicolith, presence of either pericecal or

abdominal free fluid are considered of a diagnostic fact independent from the visual status of the appendix. Patients with a history of actual pregnancy status did not undergo a computerized tomography (CT) evaluation. Although magnetic resonance imaging (MRI) is among the examination techniques for differential diagnosis of AAP in pregnant patients, none of the patients at our institution underwent an MRI procedure. Concerning macroscopic and microscopic findings, pathological examination reports are classified as appendix vermiformis (without any evidence for inflammatory cell infiltration), lymphoid hyperplasia and acute appendicitis (simple appendicitis, perforated appendicitis, gangrenous appendicitis, phlegmonous appendicitis). Rare entities of granulomatous appendicitis, fibrous obliteration, mucocele, mucinous cystadenoma and carcinoid tumor are also classified. Antibioprophylaxy is given to all patients in the two groups. Patients with intraoperative diagnosis of appendix perforation, presence of periappendicular or pelvic fluid resembling to be infected received antibiotic treatment postoperatively. Centers for Disease Control and Prevention Guideline were considered in the evaluation of postoperative surgical site infections.

### Specific Obstetric Assessment in Pre/Perioperative Period

All patients with an initial diagnosis of AAP with a reproductive age were evaluated by an obstetrics and gynecology specialist. All of the pregnant patients underwent a vaginal examination to detect any ex utero haemorrhage, miscarriage material inside vagina or cervix. Pregnant patients in second and trimester were evaluated with the transvaginal US to measure cervical longitude and detect the presence of any cervical funneling. Pregnants in a suitable gestational week were evaluated with tocography to detect presence of uterine contraction and ultrasonographic foetal nonstress test to clarify the well-being of the foetus. Pregnants in the first trimester that are with a relatively higher risk for miscarriage received supplementary progesterone. Pregnants in the third trimester with uterine contractions pointing out increased risk for preterm delivery received tocolytic treatment and supplements to promote foetal pulmonary development. All of the pregnant patients received adequate perioperative hydration to prevent dehydration.

### Statistical Analysis

The statistical analyses were performed using IBM SPSS Statistics v25.0 (Statistical Package for the Social Sciences, Inc, Chicago, IL, USA). The quantitative variables were expressed as Mean $\pm$ SD, Median and Min-Max. The qualitative variables were reported as number and percentage (%). Kolmogorov-Smirnov tests were used to assess normality distribution of quantitative variables. Mann-Whitney U test was used to compare the quantitative variables. Pearson Chi-Square and Fisher's exact tests were used to compare qualitative variables. Sensitivity, specificity, positive predictive value,

negative predictive value and accuracy value of ultrasonography in pregnant patients with presumed diagnosis of acute appendicitis were also measured. A p-value of less than 0.05 was considered statistically significant.

## RESULTS

### Pregnant versus Non-pregnant Patients Undergoing Appendectomy

A sum of 431 women in reproductive age varying from 18 to 45 underwent appendectomy with an initial diagnosis of AAP. Patients were grouped into two, according to their pregnancy status at the time of AAP diagnosis: pregnant group (n=48) and non-pregnant group (n=383). There was no statistically significant difference between the groups concerning age ( $p=0.710$ ), WBC ( $p=0.956$ ), neutrophil count ( $p=0.868$ ), lymphocyte count ( $p=0.571$ ), thrombocyte count ( $p=0.0814$ ), RDW ( $p=0.066$ ), PDW ( $p=0.183$ ), MCH ( $p=0.105$ ), MPV ( $p=0.773$ ), MCV ( $p=0.775$ ), CRP ( $p=0.0363$ ), NLR ( $p=0.486$ ), PLR ( $p=0.712$ ), PNR ( $p=0.851$ ), WLR ( $p=0.430$ ), VVNR ( $p=0.204$ ), appendix length ( $p=0.581$ ), appendix width ( $p=0.734$ ), general histopathological findings ( $p=0.580$ ), appendiceal perforation ( $p=0.204$ ) and presence of histopathological AAP ( $p=0.429$ ). On the other hand, there was a statistically significant difference between the groups for total bilirubin level ( $p=0.019$ ) and diagnosis of AAP in US evaluation ( $p=0.016$ ). Median bilirubin level was 0.58 mg/dL (mean $\pm$ SD: 0.71 $\pm$ 0.50) in the non-pregnant group, whereas it was 0.49 mg/dL (mean $\pm$ SD: 0.58 $\pm$ 0.53) in the pregnant group. US evaluation diagnosed AAP in 61% of 351 non-pregnant group, whereas this rate was 41.5% among 41 pregnant patients. Sensitivity, specificity, positive predictive value (PPV), negative predictive value (NPV) and accuracy rates for US in pregnant were 50.0%, 100.0%, 100.0%, 29.1% and 58.5% in order, AAP and appendiceal perforations for non-pregnant patients were 74% and 10.5% in order which were 79.2% and 4.2% for pregnant patients. In other words, negative appendectomy rates for non-pregnant and pregnant patients were 26% and 20.8% in order. None of the cases resulted in a surgical site or organ infection requiring either relaparotomy or percutaneous drainage. Only two cases in each group had superficial surgical site infection requiring simple drainage. Demographic, clinical and histopathological features of the two groups are given in Table 1, 2.

### Obstetric Course of Pregnant Patients Undergoing Appendectomy

A total of 13.734 deliveries were conducted at our obstetrical department during same time frame and an incidence of one case of appendectomy for preliminary diagnosis of AAP in 286 births. Also, pregnant women consisted of 11.1% of the reproductive-aged women who underwent appendectomy due to preliminary diagnosis of AAP. Among the 48 pregnant patients who underwent appendectomy, 25 (52.1%) pregnant patients were in first trimester, 14 (29.2%) pregnant patients were in second

trimester, remaining eight (16.7%) pregnant patients were in the third trimester of pregnancy. Among the 23 of the patients in the first trimester and 10 of the patients in the second trimester, appendectomy was performed using McBurney incision. Remaining two patients in the first trimester underwent laparoscopic appendectomy. Ten the second trimester patients and all of the third trimester patients underwent appendectomy via Rockey-Davis or pararectal incision, considering dimensions of the uterus and locational change of cecum. Following appendectomy, 42 (87.5%) of the pregnant patients gave vaginal birth, the remaining six (12.5%) of the pregnant patients admitted to the obstetrics clinic due to preterm delivery (3–8 weeks before gestational term). Among the preterm delivery pregnant patients, five of them gave birth via cesarean section; one pregnant patient gave vaginal birth.

Among the term deliveries, all of the babies were free of mortality and morbidity. Among the preterm delivery pregnant patients, one had diamniotic dichorionic twins and delivered two male babies on 28<sup>th</sup> week with a body weight of 1800 gr and 1900 gr. One of the preterm pregnant patients on 30<sup>th</sup> gestational week, the baby was delivered but died postpartum day 17 due to respiratory failure. Another preterm pregnancy on 22<sup>nd</sup> gestational week was terminated following the diagnosis of in-utero exitus of a 500 gr fetus. Mean birth weight of babies was 2950 gr (min-max:). As a result, 31 (64.6%) pregnant patients underwent a cesarean section, and 17 (35.4%) delivered transvaginally.

## DISCUSSION

AAP is among the most frequent non-obstetrical conditions requiring surgical treatment. Frequency of pregnant patient those diagnosed AAP and underwent appendectomy varies from 0.18 to 10.56 per every 1000 pregnant patients. Our literature analysis of 67 published studies reveals that 11.198 of 11.556.461 pregnant patients underwent surgical exploration with a preliminary diagnosis of AAP. In other words, the frequency of appendectomy per 1000 pregnant patients is 0.97 (Table 3).

Acceptable negative appendectomy (NA) is among the most frequently emphasized issues, which has a reported rate vary from 0% to 50% in many studies. The general consensus for an acceptable NA rate is 10 to 25%. However, newer studies report rates lower than 10% related to recent diagnostic instrumentation and scoring systems.<sup>[6–8]</sup> As Tubo-ovarian diseases are leading conditions resembling AAP, the NA rate is higher in female patients than in males. Literature review of the 14 published studies (n=98.933) comparing pregnant (n=3.971) and non-pregnant AAP (n=94.962) reveal that NA rates of pregnant women vary from 0% to 38% whereas NA rates of non-pregnant women vary from 0% to 21.8%.<sup>[4,5,9–20]</sup> In eight of these studies, NA was found to be higher in pregnant patients, whereas in three of these studies, NA was found to be higher in non-pregnant. Although it was not statistically significant, NA rates of pregnant patients were higher in

**Table 1.** Comparison of the pregnant and non-pregnant Acute appendicitis patients concerning clinical, biochemical and histopathological parameters

Patients' characteristics	Non-pregnant (n=383)	Pregnant (n=48)	p
	Median (min-max)	Median (min-max)	
Age	28 (18-45)	27.5 (19-45)	0.710
White blood cell	12.4 (1.2-26.9)	12.4 (6.3-22.4)	0.956
Neutrophil	9.7 (0.4-22.9)	9.6 (2.4-30.8)	0.868
Lymphocyte	1.6 (0.2-7.8)	1.7 (0.5-14.8)	0.571
Platelets	251 (43-570)	246 (141-573)	0.814
Red cell distribution width	13.7 (11-33.3)	14.5 (11.9-24.2)	0.066
Platelet distribution width	16.2 (8.4-23.9)	16.5 (9.6-19.1)	0.183
Mean corpuscular hemoglobin	28.5 (14.5-36.9)	29 (20.9-32.7)	0.105
Mean platelet volume	9 (6.1-14.5)	8.9 (5.4-12.7)	0.773
Mean corpuscular volume	84.6 (56-108)	84.9 (66-93)	0.775
Total bilirubin	0.58 (0.13-3.66)	0.49 (0.19-3.48)	0.019
C-reactive protein	1.86 (0.1-55)	2.26 (0.3-35.2)	0.363
Neutrophil to lymphocyte ratio	5.6 (0.15-41.2)	5.8 (1.7-30.8)	0.486
platelet to lymphocyte ratio	155 (22-1065)	150 (66-955)	0.712
Platelet to neutrophil ratio	25.4 (7.1-382)	25.3 (7.1-100)	0.851
White blood cell to lymphocyte ratio	7.1(1.15-46.7)	7.5 (1.1-32.2)	0.430
white blood cell to neutrophil ratio	1.3 (0.5-14.2)	1.2 (0.5-6.9)	0.204
Appendix lenght (mm)	60 (10-120)	60 (30-130)	0.581
Appendix width (mm)	10 (5-60)	10 (4-30)	0.734
	<b>n (%)</b>	<b>n (%)</b>	
Ultrasonography			0.016
Acute appendicitis (-)	137 (39)	24 (58.5)	
Acute appendicitis (+)	214 (61)	17 (41.5)	
Appendectomy type			0.001
Open	295 (77)	46 (95.8)	
Lap	88 (23)	2 (4.2)	
Histopathological findings			0.580
Appendix vermiformis	35	4	
Acute appendicitis	237	35	
Perforated acute appendicitis	40	2	
Lymphoid hyperplasia	47	3	
Carcinoid	2	1	
Mucocele	2	1	
Granulomatous appendicitis	2	0	
Fibrous obliteration	16	2	
E. Vermiculairs	2	0	
Appendiceal perforation			0.204
Yes	40 (10.5)	2 (4.2)	
No	343 (89.5)	46 (95.8)	
Acute appendicitis			0.429
Yes	283 (74)	38 (79.2)	
No	100 (26)	10 (20.8)	

**Table 2.** Comparison of the ultrasonographic assessment of pregnant and non-pregnant patients

	Pregnant (n=41)		Non- pregnant (n=351)	
	Acute appendicitis (+)	Acute appendicitis (-)	Acute appendicitis (+)	Acute appendicitis (-)
Ultrasonography acute appendicitis (+)	17	0	177	37
Ultrasonography acute appendicitis (-)	17	7	86	51
Sensitivity	50.0%		67.3%	
Specificity	100%		57.9%	
Positive predictive value	100%		82.7%	
Negative predictive value	29.2%		37.2%	
Accuracy	58.5%		65.0%	

**Table 3.** Literature review of some studies published in English language literature on ratio of the pregnant acute appendicitis

References	Journal	Total delivery or pregnancy	Pregnant appendicitis	Pregnant appendicitis/ Delivery (1000)
Aras	Rev Assoc Med Bras (1992). 2016;62:622-7	6.540	38	5.81
Masood	Obstet Gynecol Int J 2016;5: 00173	12.687	134	10.56
Aggenbach	Int J Surg. 2015;15:84-9	25.443	21	0.83
Cheng	Surg Endosc. 2015;29:1394-9	1.147.214	859	0.75
Kumamoto	Surg Today. 2015;45:1521-6	13.479	33	2.45
Abbasi	BJOG. 2014;121:1509-14	7.037.386	7114	1.01
Al- Dahamsh	J College of Med Sci-Nepal 2012; 8: 36-43	9.783	28	2.86
Jung	J Korean Soc Coloproctol 2012;28:152-9	14.203	25	1.76
Agholor	J Obstet Gynaecol Res. 2011;37:1540-8	16.173	23	1.42
Park	Eur J Obstet Gynecol Reprod Biol. 2010 ;148:44	954	8	8.39
Freeland	Am J Surg. 2009;198:753-8	65.000	23	0.35
Kazim	Int J Surg. 2009;7:365-7	43.134	37	0.86
Machado	JSLs. 2009;13:384-90	16.803	26	1.55
Zhang	Chin Med J (Engl). 2009;122:521-4	30.098	102	3.39
Al-Mulhim	Saudi J Gastroenterol. 2008;14:114-7	67.990	65	0.96
Moreno-Sanz	J Am Coll Surg. 2007;205:37-42	3.969	9	2.27
Rollins	Surg Endosc. 2004;18:237-41	18.590	30	1.61
Ueberrueck	World J Surg. 2004;28:508-11	46.969	94	2.00
Raja	Rawal Med J 2003;28:52-5	3.812	11	2.89
Eryilmaz	Dig Surg 2002;19:40-4	31.480	24	0.76
Popkin	Am J Surg 2002; 183: 20-2.	36.000	23	0.64
Duqoum	East Mediterr Health J. 2001;7:642-5	16.443	10	0.61
De Perrot	Surg Laparosc Endosc Percutan Tech.2000;10:368	3.702	9	2.43
Hoshino	Int J Gynaecol Obstet. 2000;69:271-3	15.000	15	1.00
Mourad	Am J Obstet Gynecol. 2000;182:1027-9	66.993	67	1.00
Tracey	Am Surg. 2000;66:555-9	44.845	22	0.49
Affleck	Am J Surg. 1999;178:523-9	32.818	40	1.22
Al-Qudah	J Obstet Gynaecol. 1999;19:362-4	52.108	46	0.88
Andersen	Acta Obstet Gynecol Scand. 1999;78:758-62	32.163	56	1.74

**Table 3.** Literature review of some studies published in English language literature on ratio of the pregnant acute appendicitis (continued)

References	Journal	Total delivery or pregnancy	Pregnant appendicitis	Pregnant appendicitis/ Delivery (1000)
Hee	Int J Gynaecol Obstet. 1999;65:129-35	320.949	117	0.36
Wittich	Mil Med. 1999;164:671-4	6.050	6	0.99
Al-Mulhim	Int Surg. 1996;81:295-7	31.950	52	1.63
To	Aust N Z J Surg. 1995;65:799-803	38.070	38	1.00
Lopez	J Obstetrics Gynecol 1994; 14: 133-7	41.206	62	1.50
Halvorsen	Eur J Surg. 1992;158:603-6.	44.577	16	0.36
Mahmoodian	South Med J. 1992;85:19-24	12.349	9	0.73
Al-Qasabi	Ann Saudi Med. 1991;11:58-61	31.245	46	1.47
Mazze	Obstet Gynecol. 1991;77:835-40	720.000	778	1.08
Tamir	Am J Surg. 1990;160:571-5	73.000	84	1.15
Bailey	Am Surg. 1986;52(4):218-21	100.145	41	0.41
Horowitz	Arch Surg. 1985;120:1362-7	66.351	12	0.18
Weingold	Clin Obstet Gynecol. 1983;26:801-9	19.187	24	1.25
Farquharson	Scott Med J. 1980;25:36-8	50.089	25	0.50
Punnonen	Acta Chir Scand. 1979;145:555-8	20.363	24	1.18
Gomez	Am J Surg. 1979;137:180-3	76.580	35	0.46
Babaknia	Obstet Gynecol. 1977;50:40-4	25.847	12	0.46
Zaitoon	Am Surg. 1977;43:395-8	11.844	11	0.93
Cunningham	Obstet Gynecol. 1975;45:415-20	91.800	34	0.37
Mohammed	Can Med Assoc J. 1975;112:1187-8	34.270	25	0.73
Finch	Br J Surg. 1974;61:129-32	94.000	75	0.80
Taylor	N Z J Obstet Gynaecol. 1972;12:202-3	38.719	55	1.42
O'Neill	Aust N Z J Obstet Gynaecol. 1969;9:94-9	91.500	62	0.68
Kurtz	Obstet Gynecol 1964; 23(4):528-532	84.260	41	0.49
Sarason	Obstet Gynecol. 1963;22:382-6	11.000	14	1.27
Bronstein	Am J Obstet Gynecol. 1963;86:514-6	39.000	20	0.51
King	Calif Med. 1962;97:158-62	74.000	36	0.49
Lee	JAMA. 1965;193:966-8	16.100	34	2.11
MacBeth	Can J Surg. 1961;4:419-28	59.758	50	0.84
Townsend	Am Surg. 1960;26:425-7	33.000	29	0.88
West	Am Surg. 1960;26:425-7	39.867	35	0.88
Sprong	Calif Med. 1959;91:258-60	19.932	20	1.00
Easton	Postgrad Med J. 1957;33:272-7	8.608	14	1.63
Hoffman	Am J Obstet Gynecol. 1954;67:1338-50	44.242	126	2.85
Meharg	Obstet Gynecol. 1953;1:460-5	6.106	25	4.09
Priddle	Am J Obstet Gynecol. 1951;62:150-5	59.403	51	0.86
Hamlin	N Engl J Med. 1950;244:128-31	92.772	40	0.43
Baer	JAMA. 1932;98:1359-64	16.543	28	1.69
Total	11.543.752	11.198		0.97

the study we present (26.0 vs. 20.8%). These results point out that higher NA rates demonstrate easier made surgical treatment decisions of surgeons to avoid AAP related com-

plications, such as perforation, whereas lower NA rates of pregnant demonstrate meticulously made surgical treatment decisions of surgeons or use diagnostic tools more often.

One of the subjects that AAP studies touch upon is the frequency of perforated AAP. This condition is the most important subject to be mentioned in these studies when its complications are considered. Some authors justify the relative favor between NA and complicated AAP, considering maternal and fetal complications and promote surgical treatment in all pregnant with a possibility of AAP.<sup>[20]</sup> Opponent authors emphasize the 4% rate of maternal and fetal complications among pregnant who underwent NA and be in relief against considering NA as an innocent procedure.<sup>[20,21]</sup> According to the literature analysis mentioned above, perforated AAP rates vary from 0% to 40.4% in pregnant and 3.7% to 29% in non-pregnant. Perforated AAP rate was high among pregnant in seven of these studies and was high among non-pregnant in three of these studies. These results reveal relatively high rates of both NA and perforated AAP among pregnant. In our study none of the pregnant that underwent NA experienced neither maternal nor fetal complication. On the contrary, one of the pregnant with a perforated AAP had an uncomplicated preterm delivery.

Many studies compare histopathologically proven AAP and non-AAP patient groups (control groups) to reveal sensitivity, specificity and cut-off levels of biochemical laboratory parameters (WBC, MPV, RDW, PDW, Platelets, Neutrophil, CRP, Bilirubin) in diagnosis and predicting of AAP related complications.<sup>[22,23]</sup> Similar parameters are analyzed in pregnant AAP studies as well.<sup>[24]</sup> However, a limited number of studies comparing pregnant and non-pregnant AAP patients analyse WBC and neutrophil counts, some of which reveal higher WBC or neutrophil counts in pregnant AAP group and some of them have non-significant difference.<sup>[4,1,11–15]</sup> One of the most important features of the study presented is comparing the groups concerning all the parameters mentioned above and revealing an insignificant difference between the groups except the total bilirubin levels.

Bilirubin is one of the most analyzed parameters in studies about AAP. Bacteria's hepatic involvement via portal vein following proliferation inside appendix lumen leading to the limitation in hepatic uptake and excretion and cytokine-mediated inhibition of bile salt transport have been shown previously.<sup>[24–28]</sup> Besides, hemolysis related to systemic infection has also been shown to result in an increase of bilirubin load. Many studies show a relatively higher increase in total bilirubin levels in complicated and uncomplicated AAP cases when compared with NA and the highest increase in complicated AAP.<sup>[24–28]</sup> However, some studies report a relatively increased bilirubin level among AAP group without any significant difference between complicated and non-complicated AAP groups.<sup>[28]</sup> Bilirubin level has also been shown to be predicting factor for perforation and other complications related to AAP. To our knowledge, none of the pregnant AAP studies analyzed the relation between bilirubin level and AAP. This study reveals relatively higher levels of bilirubin in the non-pregnant AAP group compared with the pregnant AAP group. Besides,

the comparison of histologically-proven AAP cases revealed relatively higher bilirubin levels in the non-pregnant group. Currently, we have no comment to clarify relatively lower bilirubin levels in the pregnant group and this study requires to be supported with further prospective studies.

We would like to share some exceptional facts of this study. First of all, non-pregnant appendectomised female patients in reproductive age were completely included in the control group to minimize the risk of bias. Secondly, the relatively lower sensitivity of US examination depends on the radiology residents on night-shift who are relatively less experienced. The third fact is the absence of an MR examination on any of the pregnant, as the MR examination and a radiology specialist were unavailable during night-shifts. The fourth is the difficulty in providing all of the patients' Alvarado scores admitted in the emergency unit of our institution despite most of the patients with a preliminary AAP diagnosis are followed with this score.

As a result, AAP is among the most frequent conditions requiring surgical treatment during pregnancy. Physiological changes of pregnant may interfere with clinical findings and biochemical parameters which leads to higher rates of perforated AAP among pregnant AAP patients. All patients with a preliminary diagnosis of AAP must be followed closely and evaluated with consecutive US examinations to minimize maternal and fetal complications.

**Ethics Committee Approval:** Approved by the local ethics committee.

**Peer-review:** Internally peer-reviewed.

**Authorship Contributions:** Concept: S.A.; Design: S.A., C.K.; Supervision: S.A., S.Y.; Fundings: C.K., B.S.; Materials: C.K., B.S., E.İ.C.; Data: C.K., B.S., E.İ.C.; Analysis: S.A.; Literature search: S.A., C.K.; Writing: S.A.; Critical revision: S.A., C.K., S.Y.

**Conflict of Interest:** None declared.

**Financial Disclosure:** The authors declared that this study has received no financial support.

## REFERENCES

1. Akbulut S, Koc C, Kocaaslan H, Gonultas F, Samdanci E, Yologlu S, et al. Comparison of clinical and histopathological features of patients who underwent incidental or emergency appendectomy. *World J Gastrointest Surg* 2019;11:19–26. [\[CrossRef\]](#)
2. Emre A, Akbulut S, Bozdog Z, Yilmaz M, Kanlioz M, Emre R, et al. Routine histopathologic examination of appendectomy specimens: retrospective analysis of 1255 patients. *Int Surg* 2013;98:354–62. [\[CrossRef\]](#)
3. Andersson RE, Lambe M. Incidence of appendicitis during pregnancy. *Int J Epidemiol* 2001;30:1281–5. [\[CrossRef\]](#)
4. Aras A, Karaman E, Pekşen Ç, Kızıltan R, Kotan MÇ. The diagnosis of acute appendicitis in pregnant versus non-pregnant women: A comparative study. *Rev Assoc Med Bras (1992)* 2016;62:622–7. [\[CrossRef\]](#)
5. Ito K, Ito H, Whang EE, Tavakkolizadeh A. Appendectomy in pregnancy: evaluation of the risks of a negative appendectomy. *Am J Surg*

- 2012;203:145–50. [CrossRef]
6. Kapan S, Bozkurt MA, Turhan AN, Gönenç M, Alış H. Management of acute appendicitis in pregnancy. *Ulus Travma Acil Cerrahi Derg* 2013;19:20–4. [CrossRef]
  7. Sammalkorpi HE, Mentula P, Savolainen H, Leppäniemi A. The Introduction of Adult Appendicitis Score Reduced Negative Appendectomy Rate. *Scand J Surg* 2017;106:196–201. [CrossRef]
  8. Schok T, Simons PC, Janssen-Heijnen ML, Peters NA, Konsten JL. Prospective evaluation of the added value of imaging within the Dutch National Diagnostic Appendicitis Guideline--do we forget our clinical eye?. *Dig Surg* 2014;31:436–43. [CrossRef]
  9. Segev L, Segev Y, Rayman S, Nissan A, Sadot E. Acute Appendicitis During Pregnancy: Different from the Nonpregnant State?. *World J Surg* 2017;41:75–81. [CrossRef]
  10. Davoodabadi A, Davoodabadi H, Akbari H, Janzamini M. Appendicitis in Pregnancy: Presentation, Management and Complications. *Zahedan J Res Med Sci* 2016;18:e7557. [CrossRef]
  11. Baruch Y, Canetti M, Blecher Y, Yogev Y, Grisaru D, Michaan N. The diagnostic accuracy of ultrasound in the diagnosis of acute appendicitis in pregnancy. *J Matern Fetal Neonatal Med* 2019;1–6. [CrossRef]
  12. Bazdar S, Dehghankhalili M, Yaghmaei S, Azadegan M, Pourdavood A, Niakan MH, et al. Acute Appendicitis during Pregnancy; Results of a Cohort Study in a Single Iranian Center. *Bull Emerg Trauma* 2018;6:122–7. [CrossRef]
  13. Bhandari TR, Shahi S, Acharya S. Acute Appendicitis in Pregnancy and the Developing World. *Int Sch Res Notices* 2017;2017:2636759. [CrossRef]
  14. Tatli F, Yucel Y, Gozeneli O, Dirican A, Uzunkoy A, Yalçın HC, et al. The Alvarado Score is accurate in pregnancy: a retrospective case-control study. *Eur J Trauma Emerg Surg* 2019;45:411–6. [CrossRef]
  15. Kumamoto K, Imaizumi H, Hokama N, Ishiguro T, Ishibashi K, Baba K, et al. Recent trend of acute appendicitis during pregnancy. *Surg Today* 2015;45:1521–6. [CrossRef]
  16. McGory ML, Zingmond DS, Tillou A, Hiatt JR, Ko CY, Cryer HM. Negative appendectomy in pregnant women is associated with a substantial risk of fetal loss. *J Am Coll Surg* 2007;205:534–40. [CrossRef]
  17. Yılmaz HG, Akgun Y, Bac B, Celik Y. Acute appendicitis in pregnancy--risk factors associated with principal outcomes: a case control study. *Int J Surg* 2007;5:192–7. [CrossRef]
  18. Finch DR, Lee E. Acute appendicitis complicating pregnancy in the Oxford region. *Br J Surg* 1974;61:129–32. [CrossRef]
  19. Richards C, Daya S. Diagnosis of acute appendicitis in pregnancy. *Can J Surg* 1989;32:358–60.
  20. Hirsch L, Yogev Y, Ashwal E, From A, Ben-Haroush A, Peled Y. The impact of pregnancy on the accuracy and delay in diagnosis of acute appendicitis. *J Matern Fetal Neonatal Med* 2014;27:1357–60. [CrossRef]
  21. Jung JY, Na JU, Han SK, Choi PC, Lee JH, Shin DH. Differential diagnoses of magnetic resonance imaging for suspected acute appendicitis in pregnant patients. *World J Emerg Med* 2018;9:26–32. [CrossRef]
  22. Boshnak N, Boshnaq M, Elgohary H. Evaluation of Platelet Indices and Red Cell Distribution Width as New Biomarkers for the Diagnosis of Acute Appendicitis. *J Invest Surg* 2018;31:121–9. [CrossRef]
  23. Ulukent SC, Sarici IS, Ulutas K. All CBC parameters in diagnosis of acute appendicitis. *Int J Clin Exp Med* 2016;9:11871–6.
  24. Çınar H, Aygün A, Derebey M, Tarım İA, Akalın Ç, Büyükkakıncak S, et al. Significance of hemogram on diagnosis of acute appendicitis during pregnancy. *Ulus Travma Acil Cerrahi Derg* 2018;24:423–8. [CrossRef]
  25. Eren T, Tombalak E, Ozemir IA, Leblebici M, Ziyade S, Ekinci O, et al. Hyperbilirubinemia as a predictive factor in acute appendicitis. *Eur J Trauma Emerg Surg* 2016;42:471–6. [CrossRef]
  26. D'Souza N, Karim D, Sunthareswaran R. Bilirubin; a diagnostic marker for appendicitis. *Int J Surg* 2013;11:1114–7. [CrossRef]
  27. Nevler A, Berger Y, Rabinovitz A, Zmora O, Shabtai M, Rosin D, et al. Diagnostic Value of Serum Bilirubin and Liver Enzyme Levels in Acute Appendicitis. *Isr Med Assoc J* 2018;20:176–81.
  28. Sushruth S, Vijayakumar C, Srinivasan K, Raj Kumar N, Balasubramanian G, Verma SK, et al. Role of C-Reactive Protein, White Blood Cell Counts, Bilirubin Levels, and Imaging in the Diagnosis of Acute Appendicitis as a Cause of Right Iliac Fossa Pain. *Cureus* 2018;10:e2070.

## ORIJİNAL ÇALIŞMA - ÖZET

### Apendektomi yapılan gebe ve gebe olmayan hastaların demografik ve klinik özelliklerinin karşılaştırılması

Dr. Cemalettin Koç,<sup>1</sup> Dr. Sami Akbulut,<sup>1</sup> Dr. Ebru İnci Coşkun,<sup>2</sup> Dr. Barış Sarıcı,<sup>1</sup> Dr. Sezai Yılmaz<sup>1</sup>

<sup>1</sup>İnönü Üniversitesi Tıp Fakültesi, Cerrahi ve Karaciğer Nakli Enstitüsü, Malatya

<sup>2</sup>İnönü Üniversitesi Tıp Fakültesi, Kadın Hastalıkları ve Doğum Anabilim Dalı, Malatya

**AMAÇ:** Bu retrospektif çalışma akut apandisit ön tanısıyla apendektomi yapılan gebe ve gebe olmayan hastaların demografik ve klinikopatolojik özelliklerini karşılaştırmayı amaçlamaktadır.

**GEREÇ VE YÖNTEM:** Haziran 2009 ve Ocak 2019 tarihleri arasında üreme çağındaki 431 kadın hastaya akut apandisit ön tanısıyla apendektomi yapıldı. Hastalar gebelik durumları gözönünde bulundurularak iki gruba ayrıldı: Gebe grup (n=48) ve gebe olmayan grup (n=383). Her iki grup demografik, klinik ve histopatolojik özellikler yönünden karşılaştırıldı.

**BULGULAR:** Gebe ve gebe olmayan gruplar arasında total bilirubin (p=0.019) ve ultrasonografik bulgular (p=0.016) dışında istatistiksel olarak anlamlı farklılık saptanmadı. Gebe olmayan grupta negatif apendektomi ve perforasyon oranları sırasıyla %26 ve %10.5 olarak hesaplanırken gebe grupta bu oranlar sırasıyla %20.8 ve %4.2 olarak hesaplandı. Ultrasonografinin gebe grubundaki sensitivite, spesifite ve doğruluk oranları sırasıyla %50, %100 ve %58.5 olarak saptanırken gebe olmayan grupta bu oranlar sırasıyla %67.3, %57.9 ve %65 olarak bulundu. Gebelerin %52.1'i birinci trimesterde, %29.2'si ikinci trimesterde ve geriye kalan %16.7'si üçüncü trimesterdeydi. Miadında gerçekleşen doğumların (%87.5) hiçbirinde fetal veya maternal komplikasyon gelişmedi. Buna karşın preterm gerçekleşen doğumların (%12.5) ikisi neonatal mortalite ile sonuçlandı.

**TARTIŞMA:** İstatistiksel olarak anlamlılık göstermemekle birlikte gebelerde negatif apendektomi ve perforasyon oranları gebe olmayan hastalara göre daha düşük bulundu ki bu durumun en önemli sebebi akut karın ile başvuran gebelerin klinik olarak daha hassas bir şekilde değerlendirilmesidir.

**Anahtar sözcükler:** Akut apandisit; apendektomi; gebelik; obstetrik komplikasyonlar; preterm eylem.

Ulus Travma Acil Cerrahi Derg 2020;26(4):555-562 doi: 10.14744/tjtes.2020.12544