

- Wollmann CG. Reel Syndrome the Ratchet mechanism. *Minerva Cardioangiol* 2011; 59: 197-202.
- Patel MB, Pandya K, Shah AJ, Lojewski E, Castellani MD, Thakur R. Reel syndrome-not a twiddler variant. *J Interv Card Electrophysiol* 2008; 23: 243-6. [CrossRef]
- Carnero -Varo A, Pérez-Paredes M, Ruiz-Ros JA, Giménez-Cervantes D, Martínez-Corbalán FR, Cubero-Lopez T, et al. "Reel Syndrome": a new form of Twiddler's syndrome? *Circulation* 1999; 100: e45-6. [CrossRef]
- Vural A, Ağaçdiken A, Ural D, Komsuoğlu B. Reel syndrome and pulsatile liver in a patient with a two-chamber pacemaker. *Jpn Heart J* 2004; 45: 1037-42. [CrossRef]
- Fyke FE, McCearley SS. Parameter signature of a reel problem. *Pacing Clin Electrophysiol* 2011; 34: 1031-3. [CrossRef]
- Ejima K, Shoda M, Manaka T, Hagiwara N. Reel syndrome. *J Cardiovasc Electrophysiol* 2009; 20: 822. [CrossRef]
- Bayliss CE, Beanlands DS, Baird RJ. The pacemaker-twiddler's syndrome: a new complication of implantable transvenous pacemakers. *Can Med Assoc J* 1968; 99: 371-3.
- Tegtmeyer CJ, Deignan JM. The cardiac pacemaker: a different twist. *AJR Am J Roentgenol* 1976; 126: 1017-8.

Address for Correspondence/Yazışma Adresi: Dr. Serkan Saygı
Çanakkale Onsekiz Mart Üniversitesi, Araştırma ve Uygulama
Hastanesi, Kardiyoloji Anabilim Dalı, 17110 Kepez, Çanakkale-Türkiye
Phone: +90 286 263 59 50 Fax: +90 286 263 59 56
E-mail: serkankard@yahoo.com, serkankard@gmail.com
Available Online Date/Çevrimiçi Yayın Tarihi: 05.11.2012



©Telif Hakkı 2013 AVES Yayıncılık Ltd. Şti. - Makale metnine www.anakarder.com web sayfasından ulaşılabilir.
©Copyright 2013 by AVES Yayıncılık Ltd. - Available on-line at www.anakarder.com
doi:10.5152/akd.2013.010

Successful percutaneous balloon mitral valvuloplasty in patients with left atrial appendage thrombus

Sol atriyal apendikte trombüs olan iki hastada başarılı mitral balon valvüloplasti

Introduction

Percutaneous balloon mitral valvuloplasty (PBMV) has become the treatment of choice for patients with symptomatic mitral stenosis since its successful use by Inoue et al. (1) in 1984. One of contraindications to this technique is the presence of thrombus in the left atrium (LA) or left atrial appendage (LAA). Nevertheless, there are some publications indicating that Inoue technique can be safely performed in patients with LAA thrombus (2).

In this report, we describe two patients with rheumatic mitral stenosis referred for PBMV and were found to have LAA thrombus. Despite the presence of LAA thrombus, successful PBMV, with the help of transthoracic echocardiography (TTE), was undertaken without complications.

Case Reports

Case 1

A 51-year-old woman was diagnosed with rheumatic mitral stenosis and referred to our hospital for PBMV. TTE revealed mild mitral regurgitation, moderate-severe mitral stenosis with a mitral valve area (MVA) of 1.1 cm² and systolic pulmonary artery pressure (PAP) of 70 mmHg.




Figure 1. Transesophageal echocardiography: a vertical plane from mid-esophagus demonstrates left atrial appendage and thrombus within the left atrial appendage (*) (Case 1)

LA - left atrium, LUPV - left upper pulmonary vein, LV - left ventricle




Figure 2. Transthoracic apical four-chamber view recorded during PBMV (Case 1)

LA - left atrium, LV - left ventricle, PBMV - percutaneous balloon mitral valvuloplasty
RA - right atrium, RV - right ventricle

Maximum and mean gradients across the valve were 23 and 11 mmHg respectively. Transesophageal echocardiography (TEE) revealed thrombus in the LAA (Fig. 1). Wilkins mitral valve score was calculated as 7. Mitral valve replacement (MVR) was offered to the patient, but she refused. PBMV was explained to the patient with risks of complications. After her informed consent for the procedure, along with TTE guidance, interatrial septum was punctured from more basal than usual and dilatation was performed by Inoue balloon with as less manipulation as possible (Fig. 2). The catheter equipment was kept at the mid LA level and away from the appendage. When the balloon was deflated, great caution was exercised to avoid the catheter tip springing up to the appendage. The procedure was completed successfully without complications. TTE showed reduction of valve gradients, maximum gradient was 8.5 mmHg and mean gradient was 4 mmHg with MVA of 1.8 cm². Systolic PAP was 30 mmHg.

Case 2

A 56-year-old woman was diagnosed with mitral stenosis and atrial fibrillation in 2007. She was being followed on β -blocker and anticoagulant therapy. She was admitted to our clinic with progressive dyspnea, which limited her daily activity. On TTE, biatrial dilatation, moderate mitral

stenosis, mild mitral regurgitation, MVA was calculated as 1.2 cm² with planimetry and 1.35 cm² with Doppler. The maximum and mean gradients across the mitral valve were 19 and 11 mmHg respectively. TEE revealed a thrombus at LAA base, not protruding into LA and measured as 1.2x1.5 cm. The Wilkins valve score was calculated as 9 (Fig. 3). The patient refused to undergo open-heart surgery for mitral valve replacement. Based on the success of the previous case, and the same type of thrombus, which was restricted to the base of the LAA, PMBV was offered and the risk of the procedure was explained in detail. PMBV was performed with the help of TTE. After completion of the procedure without any complication, echocardiographic parameters were as follows, MVA was 1.7cm², systolic PAP was 35 mmHg, maximum and mean gradients across the valve were 10 mmHg and 5 mmHg respectively.

Discussion

PBMV is the treatment of choice for patients with rheumatic mitral stenosis and suitable valve anatomy. Stroke was one of the catastrophic complications of PBMV. While ACC/AHA guidelines emphasizes LA thrombus as a contraindication for PBMV (level of evidence C), there is no solid evidence for cases with left atrial appendage thrombus (3).

Some operators do not consider LAA thrombus as an absolute contraindication (4).

Koca et al. (5) performed PBMV to nine patients with symptomatic mitral stenosis and thrombus restricted to the LAA. These procedures were performed under TEE guidance and there were no thromboembolic events. Consequently, researchers concluded that in selected cases, PBMV under TEE guidance was safe and thrombus restricted to the LAA is not an absolute contraindication to the procedure.

The most comprehensive research about this comes from small scale trials comprising 28 and 30 patients. Manjunath et al. (4) performed PBMV to 30 patients with LAA thrombus and observed no systemic thromboembolic event. In this study, Manjunath et al. (4) grouped LA thrombus into five subtypes. In patients with type 1a (LAA thrombus confined to appendage), type 1b (LA appendage thrombus protruding into LA cavity) and type 2a (LA roof thrombus limited to a plane above the plane of fossa ovalis) thrombus, PBMV was considered as safe and effective with modified techniques. Shaw et al. (6) performed PBMV to 28 patients with LAA thrombus and none of the patients experienced embolic event.

Our patients had type 1a thrombus according to the classification denoted by Manjunath et al. (4). The procedures were performed with lower septal puncture and with less manipulation. Contrary to previous researchers, we used TTE.



Figure 3. Transesophageal echocardiography shows LAA thrombus at mid-esophageal aortic valve short-axis view (Case 2). Arrow-thrombus in LAA

LA - left atrium, LAA - left atrial appendage, RA - right atrium

Conclusion

PBMV is a safe option for patients with suitable valve anatomy and thrombus localized to LAA. Systemic thromboembolism is rare if performed by an experienced operator. TTE seems as a safe and effective alternative to TEE.

Hakan Akıllı, Alp Arıbaş, Gökhan Altunbaş, Kurtuluş Özdemir
Department of Cardiology, Necmettin Erbakan University, Meram
Faculty of Medicine, Konya-Turkey

References

1. Inoue K, Owaki T, Nakamura T, Kitamura F, Miyamoto N. Clinical application of transvenous mitral commissurotomy by a new balloon catheter. J Thorac Cardiovasc Surg 1984; 87: 394-402.
2. Tessier P, Mercier LA, Burelle D, Bonan R. Results of percutaneous mitral commissurotomy in patients with a left atrial appendage thrombus detected by transesophageal echocardiography. J Am Soc Echocardiogr 1994; 7: 394-9.
3. Bonow RO, Carabello BA, Chatterjee K, de Leon AC Jr, Faxon DP, Freed MD, et al. 2008 focused update incorporated into the ACC/AHA 2006 guidelines for the management of patients with valvular heart disease: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to revise the 1998 guidelines for the management of patients with valvular heart disease). Endorsed by the Society of Cardiovascular Anesthesiologists, Society for Cardiovascular Angiography and Interventions, and Society of Thoracic Surgeons. J Am Coll Cardiol 2008; 52: e1-142. [CrossRef]
4. Manjunath CN, Srinivasa KH, Ravindranath KS, Manohar JS, Prabhavathi B, Dattatreya PV, et al. Balloon mitral valvotomy in patients with mitral stenosis and left atrial thrombus. Catheter Cardiovasc Interv 2009; 74: 653-61. [CrossRef]
5. Koca V, Bozat T, Yavuz Ş, Özdemir A. Sol atriyal apendiks trombüsü olan romatizmal mitral darlığı olgularında perkütan mitral balon valvülotomi. Turk Kardiyol Dern Arş 2000; 28: 302-5.
6. Shaw TR, Northridge DB, Sutaria N. Mitral balloon valvotomy and left atrial thrombus. Heart 2005; 91: 1088-9. [CrossRef]

Address for Correspondence/Yazışma Adresi: Dr. Hakan Akıllı
Necmettin Erbakan Üniversitesi Meram Tıp Fakültesi, Kardiyoloji
Sekreterliği, Meram, 42090 Konya-Türkiye
Phone: +90 332 223 79 41 Fax: +90 332 223 68 81
E-mail: hakanakilli@hotmail.com

Available Online Date/Çevrimiçi Yayın Tarihi: 05.11.2012

©Telif Hakkı 2013 AVES Yayıncılık Ltd. Şti. - Makale metnine www.anakarder.com web sayfasından ulaşılabilir.

©Copyright 2013 by AVES Yayıncılık Ltd. - Available on-line at www.anakarder.com
doi:10.5152/akd.2013.011



The role of two-dimensional speckle-tracking echocardiography in a patient with Behçet's disease

Behçet hastalığı olan hastada iki boyutlu benek takip yöntemi ekokardiyografinin önemi

Introduction

Behçet's disease (BD) is a systemic inflammatory disorder of unknown origin characterized by variable clinical manifestations. Most