

# Brain Abscess due to *Streptococcus Intermedius* Secondary to Tetralogy of Fallot in a Child: A Case Report

## Fallot Tetralojisine Sekonder Gelişen *Streptococcus Intermedius*'un Neden Olduğu Beyin Apresi: Olgu Sunumu

Esra ÇELİK KUZAYTEPE,<sup>1</sup> Ayşe KARAASLAN,<sup>1</sup> Yasemin AKIN,<sup>1</sup> Tufan HİÇDÖNMEZ,<sup>2</sup> Ömer ÇİFTÇİ,<sup>3</sup> İbrahim MERİÇ,<sup>1</sup> Esmâ ESMİ,<sup>1</sup> Melis ŞİRİNOĞLU,<sup>1</sup> Serap GENÇ YÜZÜAK,<sup>1</sup> Şerife DÜLGER<sup>1</sup>

<sup>1</sup>Department of Pediatrics, Kartal Dr. Lütfi Kırdar Training and Research Hospital, İstanbul, Turkey

<sup>2</sup>Department of Neurosurgery, Kartal Dr. Lütfi Kırdar Training and Research Hospital, İstanbul, Turkey

<sup>3</sup>Department of Pediatric Cardiology, Koşuyolu Yüksek İhtisas Training and Research Hospital, İstanbul, Turkey

### Summary

A 10-year-old Syrian boy was admitted to emergency room with complaints of headache, vomiting, decreased level of consciousness, and imbalance when walking. On physical examination, clubbing of fingers and perioral cyanosis were observed. Cranial computed tomography examination revealed 41.55x25.65 mm abscess formation in right occipitoparietal region. Abscess aspiration was performed. Empirical intravenous ceftriaxone and metronidazole therapy was initiated. Abscess culture yielded *Streptococcus intermedius*; however, urine culture yielded extended-spectrum beta-lactamase-producing *Escherichia coli*, and in 2 successive blood cultures, methicillin-resistant coagulase-negative staphylococci were found. Therefore, antibiotic therapy was replaced with IV vancomycin and meropenem combination. Follow-up magnetic resonance imaging was performed and abscess formulation was observed. Total abscess excision was performed. Patient was successfully treated with surgical drainage and 6 weeks of systemic antibiotic therapy. *S. intermedius* is found in normal oral flora, but can become pathogenic and cause potentially lethal infection in presence of underlying disease, such as cyanotic heart disease.

**Keywords:** Brain abscess; children; tetralogy of Fallot.

### Özet

Fallot tetralojisi tanı 10 yaşında Suriyeli erkek hasta acil servise baş ağrısı, kusma, bilinç bulanıklığı ve dengesiz yürüme şikayetleriyle başvurdu. Fizik muayenede, parmaklarda belirgin çomaklaşma ve peroral siyanoz gözlemlendi. Hastanın yapılan kranial bilgisayarlı tomografisinde sağ oksipitoparietal bölgede 41.55x25.65 mm büyüklüğünde apse formasyonu saptandı. Aps aspirasyonu gerçekleştirildi. Ampirik olarak hastaya intravenöz (iv) seftriakson ve metronidazol tedavileri başlandı. Apsenin kültüründe *Streptococcus intermedius*, idrar kültüründe genişlemiş spektrumlu bata-laktaaz üreten *Escherichia coli* ve 2 ardışık kan kültürlerinde metisiline dirençli koagülaz-negatif stafilokoklar saptandı. Bu nedenle mevcut antibiyoterapi yerine iv vankomisin ve meropenem kombinasyonu kullanıldı. Takip sırasında manyetik rezonans görüntüleme yapıldı ve apse oluşumu gözlemlendi. Aps tümüyle eksize edildi. Hasta cerrahi drenaj ve 6 hafta sistemik antibiyotik tedavisiyle başarıyla tedavi edildi. Normal ağız florasında *S. intermedius* bulunmuş olup siyanotik kalp hastalığı gibi altta yatan bir hastalık varlığında *S. intermedius* patojenik hale gelebilir ve potansiyel olarak ölümcül enfeksiyona neden olabilir.

**Anahtar sözcükler:** Beyin apresi; çocuk; Fallot tetralojisi.

**Correspondence:** Dr. Yasemin Akın.  
Kartal Dr. Lütfi Kırdar Eğitim ve Araştırma Hast.,  
Çocuk Sağ. ve Hast. Kliniği, Kartal, İstanbul, Turkey  
**Tel:** +90 216 - 441 39 00 / 2534

**Received:** 16.06.2016  
**Accepted:** 28.07.2016  
**Online edition:** 15.12.2016  
**e-mail:** jasminetr@hotmail.com



## Introduction

Brain abscess is infectious condition that causes focal suppurative collection in the brain parenchyma. It may be seen in children at every age; however, it is most frequently encountered in children aged between 4 and 7 years.<sup>[1]</sup> It is rare, but when observed, it induces life-threatening clinical condition. With development of imaging modalities (computed tomography [CT], and subsequently magnetic resonance imaging [MRI]), and appropriate antibiotherapies, treatment success rates have increased. However, while morbidity and mortality rates have decreased, it is still an important cause of mortality.<sup>[2]</sup> Brain abscess may develop as outcome of congenital heart disease, meningitis, mastoiditis, orbital cellulitis, intraoral infection, surgical intervention, or penetrating head trauma, but in 15% to 30% of cases, etiology cannot be determined.<sup>[3]</sup> Prevalence of brain abscess in congenital cyanotic heart disease ranges between 6% and 51%.<sup>[4,5]</sup> In 30% to 34% of patients with brain abscess, an underlying heart defect was seen.<sup>[6]</sup> Tetralogy of Fallot (TOF) is among the most frequently seen cyanotic heart diseases that can contribute to brain abscess.

Clinical picture of children with brain abscess changes depending on size and location of abscess, presence of edema around abscess, and virulence of infectious agent, if infectious etiology is found. Most frequently observed symptoms are fever, headache, seizures, vomiting, and impaired consciousness.<sup>[7]</sup>

*Streptococcus intermedius* belongs to the *Viridans streptococci* group, found in normal oral and gastrointestinal flora. However, it can cause serious infectious conditions in abscess. Presently described is case of pediatric patient with diagnosis of TOF who developed brain abscess secondary to *S. intermedius* infection.

## Case Report

A 10-year-old Syrian boy was brought to pediatric intensive care unit with complaints of headache, vomiting, clouded consciousness, and unsteady gait. Headache had persisted for 10 days. On physical examination, subfebrile body temperature, marked clubbing of fingers, perioral cyanosis, ecchymotic lips and gingiva, and grade 3/6 pansystolic murmur were detected. Some of his biochemical parameters were as follows: white blood cell count: 35400/mm<sup>3</sup>, C-reactive protein: 187 mg/L, and sedimentation rate: 23 mm/hr. Blood and urine cultures were obtained before lumbar

puncture cranial imaging scheduled as result of detection of neurological symptoms. Cranial CT revealed lesion measuring 41.55x25.65 mm in right occipitoparietal region, consistent with abscess formation (Figure 1). Patient underwent emergency abscess aspiration in department of neurosurgery, and empirical intravenous (IV) antibiotherapy with ceftriaxone and metronidazole was initiated. *S. intermedius* was observed on culture of abscess material. Based on result of antibacterial susceptibility tests, maintenance of IV antibiotherapy with ceftriaxone and metronidazole was planned. However, because urine culture was positive for extended-spectrum beta-lactamase-producing *Escherichia coli*, and on 2 occasions blood culture was positive for methicillin-resistant coagulase-negative staphylococci, IV antibiotherapy with vancomycin and meropenem was initiated. Department of pediatric cardiology was consulted and patient was diagnosed as incomplete TOF. It was learned that patient had not had recommended cardiac surgery in Syria and had not attended follow-up cardiology visits. Our facility took over his cardiac treatment. Echocardiographic examinations were repeated 3 times and no evidence suggesting infective endocarditis was detected. Cranial MRI repeated at third week of treatment revealed minimal regression of abscess formation, so total excision of abscess was performed by department of neurosurgery. Brain abscess material was sent for culture



**Figure 1.** Brain abscess as observed in cranial computed tomography. Colored images can be seen in online issue of the journal ([www.kehadergi.com](http://www.kehadergi.com)).

a second time, and no bacterial growth was observed. Brain abscess was successfully treated with surgical intervention and systemic antibiotherapy lasting for 6 weeks. Patient was discharged and his further cardiac examinations and treatment were planned. Written informed consent was obtained from the patient who participated in this study.

## Discussion

Brain abscess is a rarely seen but life-threatening disease of childhood. Since the 1990s, due to development of additional diagnostic and therapeutic capabilities, prognosis has improved considerably. However, neurological deficits can be still be seen after treatment, and in some children, ventriculoperitoneal shunt may be required.

Brain abscess is an intracerebral focal infection that develops due to spread of local infection. Generally, purulent collection is seen in well-vascularized capsule.<sup>[8]</sup> Location of abscess typically depends on source of infection. In present patient, in presence of congenital heart disease, parenchymal tissue of parietal, and occipital regions were involved.<sup>[9]</sup> In cyanotic heart disease, bacteria cannot be filtered through pulmonary vascular bed and may spread to systemic circulation.<sup>[10]</sup> In our case, presence of TOF as underlying cyanotic heart disease and isolation of *S. intermedius* on culture of abscess material suggest this phenomenon as probable etiology.

The most frequently detected bacterial agents of brain abscess in children are aerobic and anaerobic streptococci (60–70%), gram-negative anaerobic bacilli (20–40%), *Enterobacteriaceae* (20–30%), *S. aureus* (10–15%), and fungi (1–5%).<sup>[11,12]</sup>

As a member of *S. anginosus* family, *S. intermedius* is a gram-positive catalase-negative coccus. It generally tends to cause infectious conditions, including abscess formation, generally seen in oral cavity, head and neck region, and abdomen. All members of *S. anginosus* family produce pyrogenic exotoxin. As a distinct feature, *S. intermedius* produces cytolytic enzyme intermedilysin, which is thought to facilitate formation of abscess localized in liver and deep anatomical structures.<sup>[13]</sup> Previous studies have also shown that this group of bacteria can cause development of brain abscess.<sup>[14–16]</sup> Access to central nervous system through oral, dental, or gastrointestinal tract as result of bacteremia has been suggested. As seen in

our patient, cardiac etiologies can be eliminated with echocardiographic examination. In the treatment of brain abscess caused by members of *S. anginosus* family, surgical drainage of abscess and IV administration of ceftriaxone is a good treatment alternative. Concomitant use of metronidazole for synchronous anaerobic infection has also been recommended.<sup>[17]</sup> IV treatment should be maintained for 4 to 6 weeks. Afterward, patient should be followed-up monthly for at least 3 months using imaging modalities to evaluate treatment response.<sup>[18]</sup> In present case, treatment was completed in 6 weeks; however, patient was lost to follow-up.

In conclusion, *S. intermedius*, a bacterium found in normal oral flora, can become a pathogenic agent in the presence of underlying facilitatory factors, such as TOF, which can lead to development of life-threatening infection, including brain abscess.

## Conflict of interest

None declared.

## References

1. Kaplan K. Brain abscess. *Med Clin North Am* 1985;69:345–60. [Crossref](#)
2. Mises J, Daviet F, Moussalli-Salefranque F, Sternberg B, Flandin C, Renier D. Brain abscess in the newborn infant (27 cases: initial electroclinical study, course. *Rev Electroencephalogr Neurophysiol Clin* 1987;17:301–8. [Crossref](#)
3. Wispelwey B, Scheld WM. Brain abscess. In: Mandell GL, Bennett JE, Dolin R, editors. *Principles and Practice of Infectious Diseases*. 4th ed. New York: Churchill Livingstone; 1995. p. 887–900.
4. Kumar K. Neurological complications of congenital heart disease. *Indian J Pediatr* 2000;67:287–91. [Crossref](#)
5. Frazier JL, Ahn ES, Jallo GI. Management of brain abscesses in children. *Neurosurg Focus* 2008;24:E8. [Crossref](#)
6. Agarwal A, Gergits F 3rd, Isaacson G. Metastatic intracranial abscesses of bronchopulmonary origin. *Pediatr Infect Dis J* 2003;22:277–80. [Crossref](#)
7. Wong TT, Lee LS, Wang HS, Shen EY, Jaw WC, Chiang CH, et al. Brain abscesses in children—a cooperative study of 83 cases. *Childs Nerv Syst* 1989;5:19–24. [Crossref](#)
8. Amano K, Kamano S. Cerebellar abscess due to penetrating orbital wound. *J Comput Assist Tomogr* 1982;6:1163–6. [Crossref](#)
9. Sáez-Llorens XJ, Umaña MA, Odio CM, McCracken GH Jr, Nelson JD. Brain abscess in infants and children. *Pediatr Infect Dis J* 1989;8:449–58. [Crossref](#)
10. Cochrane DD. Consultation with the specialist. *Brain abscess*. *Pediatr Rev* 1999;20:209–15. [Crossref](#)

11. Barlas O, Sencer A, Erkan K, Eraksoy H, Sencer S, Bayindir C. Stereotactic surgery in the management of brain abscess. *Surg Neurol* 1999;52:404–11. [Crossref](#)
12. Brouwer MC, Coutinho JM, van de Beek D. Clinical characteristics and outcome of brain abscess: systematic review and meta-analysis. *Neurology* 2014;82:806–13.
13. Nagamune H, Whiley RA, Goto T, Inai Y, Maeda T, Hardie JM, et al. Distribution of the intermedilysin gene among the anginosus group streptococci and correlation between intermedilysin production and deep-seated infection with *Streptococcus intermedius*. *J Clin Microbiol* 2000;38:220–6.
14. Greenlee JE. Subdural Empyema. *Curr Treat Options Neurol* 2003;5:13–22. [Crossref](#)
15. Petti CA, Simmon KE, Bender J, Blaschke A, Webster KA, Conneely MF, et al. Culture-Negative intracerebral abscesses in children and adolescents from *Streptococcus anginosus* group infection: a case series. *Clin Infect Dis* 2008;46:1578–80. [Crossref](#)
16. Felsenstein S, Williams B, Shingadia D, Coxon L, Riordan A, Demetriades AK, et al. Clinical and microbiologic features guiding treatment recommendations for brain abscesses in children. *Pediatr Infect Dis J* 2013;32:129–35. [Crossref](#)
17. Kowlessar PI, O'Connell NH, Mitchell RD, Elliott S, Elliott TS. Management of patients with *Streptococcus milleri* brain abscesses. *J Infect* 2006;52:443–50. [Crossref](#)
18. Ziai WC, Lewin JJ 3rd. Update in the diagnosis and management of central nervous system infections. *Neurol Clin* 2008;26:427–68. [Crossref](#)